



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/14, 15/29, 15/53, 15/82	A1	(11) International Publication Number: WO 96/10075 (43) International Publication Date: 4 April 1996 (04.04.96)
(21) International Application Number: PCT/US95/11855 (22) International Filing Date: 25 September 1995 (25.09.95) (30) Priority Data: 08/314,596 26 September 1994 (26.09.94) US 08/320,982 11 October 1994 (11.10.94) US Not furnished 20 September 1995 (20.09.95) US (71)(72) Applicants and Inventors: SOMERVILLE, Chris [US/US]; 5 Valley Oak, Portola Valley, CA 94028 (US). BROWN, Pierre [FR/US]; 1249 Capuchino, Burlingame, CA 94010 (US). VAN DE LOO, Frank, J. [AU/US]; 224 Dantzler Court, Lexington, KY 40503 (US). (74) Agents: KOKULIS, Paul, N. et al.; Cushman, Darby & Cushman, L.L.P., 1100 New York Avenue, N.W., Washington, DC 20005 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PRODUCTION OF HYDROXYLATED FATTY ACIDS IN GENETICALLY MODIFIED PLANTS (57) Abstract This invention relates to plant fatty acyl hydroxylases. Methods to use conserved amino acid or nucleotide sequences to obtain plant fatty acyl hydroxylases are described. Also described is the use of cDNA clones encoding a plant hydroxylase to produce a family of hydroxylated fatty acids in transgenic plants.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**PRODUCTION OF HYDROXYLATED FATTY ACIDS
IN GENETICALLY MODIFIED PLANTS**

TECHNICAL FIELD

The present invention concerns the identification of nucleic acid sequences and constructs, and methods related thereto, and the use of these sequences and constructs to produce genetically modified plants for the purpose of altering the fatty acid composition of plant oils, waxes and related compounds.

DEFINITIONS

The subject of this invention is a class of enzymes that

introduce a hydroxyl group into several different fatty acids resulting in the production of several different kinds of hydroxylated fatty acids. In particular, these enzymes catalyze hydroxylation of oleic acid to 12-hydroxy oleic acid and icosenoic acid to 14-hydroxy icosenoic acid. Other fatty acids such as palmitoleic and erucic acids may also be substrates. Since it is not possible to refer to the enzyme by reference to a unique substrate or product, we refer to the enzyme throughout as kappa hydroxylase to indicate that the enzyme introduces the hydroxyl three carbons distal (i.e., away from the carboxyl carbon of the acyl chain) from a double bond located near the center of the acyl chain.

The following fatty acids are also the subject of this invention: ricinoleic acid, 12-hydroxyoctadec-cis-9-enoic acid (12OH-18:1^{cisΔ⁹}); lesquerolic acid, 14-hydroxy-cis-11-icosenoic acid (14OH-20:1^{cisΔ¹¹}); densipolic acid, 12-hydroxyoctadec-cis-9,15-dienoic acid (12OH-18:2^{cisΔ^{9,15}}); auricolic acid, 14-hydroxy-cis-11,17-icosadienoic acid (14OH-20:2^{cisΔ^{11,17}}); hydroxyerucic, 16-hydroxydocos-cis-13-enoic acid (16OH-22:1^{cisΔ¹³}); hydroxypalmitoleic, 12-hydroxyhexadec-cis-9-enoic (12OH-16:1^{cisΔ⁹}); icosenoic acid (20:1^{cisΔ¹¹}). It will be noted that icosenoic acid is spelled eicosenoic acid in some countries.

BACKGROUND

Extensive surveys of the fatty acid composition of seed oils from different species of higher plants have resulted in the identification of at least 33 structurally distinct

monohydroxylated plant fatty acids, and 12 different polyhydroxylated fatty acids that are accumulated by one or more plant species (reviewed by van de Loo et al. 1993). Ricinoleic acid, the principal constituent of the seed oil from the castor plant *Ricinus communis* (L.), is of commercial importance. We have previously described the cloning of a gene from this species that encodes a fatty acid hydroxylase, and the use of this gene to produce ricinoleic acid in transgenic plants of other species. The scientific evidence supporting this claim was published in 1995. The use of the castor hydroxylase gene to also produce other hydroxylated fatty acids such as lesquerolic acid, densipolic acid, hydroxypalmitoleic, hydroxyerucic and auricolic acid in transgenic plants is the subject of this invention. In addition, the identification of a gene encoding a homologous hydroxylase from *Lesquerella fendleri*, and the use of this gene to produce these hydroxylated fatty acids in transgenic plants is the subject of this invention.

Castor is a minor oilseed crop. Approximately 50% of the seed weight is oil (triacylglycerol) in which 85-90% of total fatty acids are the hydroxylated fatty acid, ricinoleic acid. Oil pressed or extracted from castor seeds has many industrial uses based upon the properties endowed by the hydroxylated fatty acid. The most important uses are production of paints and varnishes, nylon-type synthetic polymers, resins, lubricants, and cosmetics (Atsmon 1989). In addition to oil,

the castor seed contains the extremely toxic protein ricin, allergenic proteins, and the alkaloid ricinine. These constituents preclude the use of the untreated seed meal (following oil extraction) as a livestock feed, normally an important economic aspect of oilseed utilization.

Furthermore, with the variable nature of castor plants and a lack of investment in breeding, castor has few favorable agronomic characteristics. For a combination of these reasons, castor is no longer grown in the United States and the development of an alternative domestic source of hydroxylated fatty acids would be attractive. The production of ricinoleic acid, the important constituent of castor oil, in an established oilseed crop through genetic engineering would be a particularly effective means of creating a domestic source.

Because there is no practical source of lesquerolic, densipolic and auricolic acids from plants that are adapted to modern agricultural practices, there is currently no large-scale use of these fatty acids by industry. However, the fatty acids would have uses similar to those of ricinoleic acid if they could be produced in large quantities at comparable cost to other plant-derived fatty acids (Smith 1985). Plant species, such as certain species in the genus *Lesquerella*, that accumulate a high proportion of these fatty acids, have not been domesticated and are not currently considered a practical source of fatty acids (Hirsinger, 1989). This invention represents a useful step toward the eventual production of these and other hydroxylated fatty

acids in transgenic plants of agricultural importance.

The taxonomic relationships between plants having similar or identical kinds of unusual fatty acids have been examined (van de Loo et al., 1993). In some cases, particular fatty acids occur mostly or solely in related taxa. In other cases there does not appear to be a direct link between taxonomic relationships and the occurrence of unusual fatty acids. In this respect, ricinoleic acid has now been identified in 12 genera from 10 families (reviewed in van de Loo et al., 1993). Thus, it appears that the ability to synthesize hydroxylated fatty acids has evolved several times independently during the radiation of the angiosperms. This suggested to us that the enzymes which introduce hydroxyl groups into fatty acids arose by minor modifications of a related enzyme. Indeed, as shown herein, the sequence similarity between $\Delta 12$ fatty acid desaturases and the kappa hydroxylase from castor is so high that it is not possible to unambiguously determine whether a particular enzyme is a desaturase or a hydroxylase on the basis of evidence in the scientific literature. Similarly, a patent application (WO (94/11516) that purports to teach the isolation and use of $\Delta 12$ fatty acid desaturases does not teach how to distinguish a hydroxylase from a desaturase. In view of the importance of being able to distinguish between these activities for the purpose of genetic engineering of plant oils, the utility of that application is limited to the several instances where direct experimental evidence (e.g., altered fatty acid composition in transgenic plants) was presented to support the assignment of function. A method for

distinguishing between fatty acid desaturases and fatty acid hydroxylases on the basis of amino acid sequence of the enzyme is also a subject of this invention.

5 A feature of hydroxylated or other unusual fatty acids is that they are generally confined to seed triacylglycerols, being largely excluded from the polar lipids by unknown mechanisms (Battey and Ohlrogge 1989; Prasad et al., 1987). This is particularly intriguing since diacylglycerol is a precursor of both triacylglycerol and polar lipid. With
10 castor microsomes, there is some evidence that the pool of ricinoleoyl-containing polar lipid is minimized by a preference of diacylglycerol acyltransferase for ricinoleate-containing diacylglycerols (Bafor et al. 1991). Analyses of vegetative tissues have generated few reports of unusual fatty
15 acids, other than those occurring in the cuticle. The cuticle contains various hydroxylated fatty acids which are interesterified to produce a high molecular weight polyester which serves a structural role. A small number of other exceptions exist in which unusual fatty acids are found in
20 tissues other than the seed.

The biosynthesis of ricinoleic acid from oleic acid in the developing endosperm of castor (*Ricinus communis*) has been studied by a variety of methods. Morris (1967) established in double-labeling studies that hydroxylation occurs directly by
25 hydroxyl substitution rather than via an unsaturated-, keto- or epoxy-intermediate. Hydroxylation using oleoyl-CoA as precursor can be demonstrated in crude preparations or microsomes, but activity in microsomes is unstable and

variable, and is lation of the microsomes involved a considerable, or sometimes complete loss of activity (Galliard and Stumpf, 1966; Moreau and Stumpf, 1981). Oleic acid can replace oleoyl-CoA as a precursor, but only in the presence of CoA, Mg^{2+} and ATP (Galliard and Stumpf, 1966) indicating that activation to the acyl-CoA is necessary. However, no radioactivity could be detected in ricinoleoyl-CoA (Moreau and Stumpf, 1981). These and more recent observations (Bafor et al., 1991) have been interpreted as evidence that the substrate for the castor oleate hydroxylase is oleic acid esterified to phosphatidylcholine or another phospholipid.

The hydroxylase is sensitive to cyanide and azide, and dialysis against metal chelators reduces activity, which could be restored by addition of $FeSO_4$, suggesting iron involvement in enzyme activity (Galliard and Stumpf, 1966). Ricinoleic acid synthesis requires molecular oxygen (Galliard and Stumpf, 1966; Moreau and Stumpf 1981) and requires NAD(P)H to reduce cytochrome b5 which is thought to be the intermediate electron donor for the hydroxylase reaction (Smith et al., 1992).

Carbon monoxide does not inhibit hydroxylation, indicating that a cytochrome P450 is not involved (Galliard and Stumpf, 1966; Moreau and Stumpf 1981). Data from a study of the substrate specificity of the hydroxylase show that all substrate parameters (i.e., chain length and double bond position with respect to both ends) are important; deviations in these parameters caused reduced activity relative to oleic acid (Howling et al., 1972). The position at which the hydroxyl was introduced, however, was determined by the

position of the double bond, always being three carbons distal. Thus, the castor acyl hydroxylase enzyme can produce a family of different hydroxylated fatty acids depending on the availability of substrates. Thus, as a matter of convenience, we refer to the enzyme throughout as a kappa hydroxylase (rather than an oleate hydroxylase) to indicate the broad substrate specificity.

The castor kappa hydroxylase has many superficial similarities to the microsomal fatty acyl desaturases (Browse and Somerville, 1991). In particular, plants have a microsomal oleate desaturase active at the $\Delta 12$ position. The substrate of this enzyme (Schmidt et al., 1993) and of the hydroxylase (Bafor et al., 1991) appears to be a fatty acid esterified to the *sn*-2 position of phosphatidylcholine. When oleate is the substrate, the modification occurs at the same position ($\Delta 12$) in the carbon chain, and requires the same cofactors, namely electrons from NADH via cytochrome *b₅* and molecular oxygen. Neither enzyme is inhibited by carbon monoxide (Moreau and Stumpf, 1981), the characteristic inhibitor of cytochrome P450 enzymes.

There do not appear to have been any published biochemical studies of the properties of the hydroxylase enzyme(s) in *Lesquerella*.

Conceptual basis of the invention

An earlier application has described the use of a cDNA clone from castor for the production of ricinoleic acid in transgenic plants. As noted above,

biochemical studies by others had suggested that the castor hydroxylase may not have strict specificity for oleic acid but would also catalyze hydroxylation of other fatty acids such as icosenoic acid (20:1^{cisΔ11}) (Howling et al., 1972). Based on these studies,

the expression of the castor hydroxylase in transgenic plants of species such as *Brassica napus* and *Arabidopsis thaliana* that accumulate fatty acids such as icosenoic acid (20:1^{cisΔ11}) and erucic acid (13-docosenoic acid; 22:1^{cisΔ13}) would be expected to accumulate some of the hydroxylated derivatives of these fatty acids due to the activity of the hydroxylase on these fatty acids. We have now obtained additional direct evidence for such a claim based on the production of ricinoleic, lesquerolic, densipolic and auricolic fatty acids in transgenic *Arabidopsis* plants and have included such evidence herein as Example 1.

We have previously disclosed the various methods by which the castor hydroxylase clone and sequences derived thereof could be used to identify other hydroxylase clones from plant species such as *Lesquerella fendleri* that are known to accumulate hydroxylated fatty acids in seed oils. In this continuation we have provided an example of the use of that aspect of the invention for the isolation of a novel hydroxylase gene from *Lesquerella fendleri*.

In view of the high degree of sequence similarity between Δ12 fatty acid desaturases and the castor hydroxylase (van de Loo et al., 1995), the validity of claims for the use of

desaturase or hydroxylase genes or sequences derived therefrom for the identification of genes of identical function from other species must be viewed with skepticism. In this application, we teach a method by which hydroxylase genes can be distinguished from desaturases and describe methods by which $\Delta 12$ desaturases can be converted to hydroxylases by the modification of the gene encoding the desaturases. A mechanistic basis for the similar reaction mechanisms of desaturases and hydroxylases was presented in an earlier patent application. Briefly, the available evidence suggests that fatty acid desaturases have a similar reaction mechanism to the bacterial enzyme methane monooxygenase which catalyses a reaction involving oxygen-atom transfer ($\text{CH}_4 \rightarrow \text{CH}_3\text{OH}$) (van de Loo et al., 1993). The cofactor in the hydroxylase component of methane monooxygenase is termed a μ -oxo bridged diiron cluster (FeOFe). The two iron atoms of the FeOFe cluster are liganded by protein-derived nitrogen or oxygen atoms, and are tightly redox-coupled by the covalently-bridging oxygen atom. The FeOFe cluster accepts two electrons, reducing it to the diferrous state, before oxygen binding. Upon oxygen binding, it is likely that heterolytic cleavage also occurs, leading to a high valent oxoiron reactive species that is stabilized by resonance rearrangements possible within the tightly coupled FeOFe cluster. The stabilized high-valent oxoiron state of methane monooxygenase is capable of proton extraction from methane, followed by oxygen transfer, giving methanol. The FeOFe cofactor has been shown to be directly relevant to plant fatty

acid modifications by the demonstration that castor stearyl-ACP desaturase contains this type of cofactor (Fox et al., 1993).

On the basis of the foregoing considerations, we hypothesized that the castor oleate hydroxylase is a structurally modified fatty acyl desaturase, based upon three arguments. The first argument involves the taxonomic distribution of plants containing ricinoleic acid. Ricinoleic acid has been found in 12 genera of 10 families of higher plants (reviewed in van de Loo et al., 1993). Thus, plants in which ricinoleic acid occurs are found throughout the plant kingdom, yet close relatives of these plants do not contain the unusual fatty acid. This pattern suggests that the ability to synthesize ricinoleic acid has arisen (and been lost) several times independently, and is therefore a quite recent divergence. In other words, the ability to synthesize ricinoleic acid has evolved rapidly, suggesting that a relatively minor genetic change in the structure of the ancestral enzyme was necessary to accomplish it.

The second argument is that many biochemical properties of castor kappa hydroxylase are similar to those of the microsomal desaturases, as discussed above (e.g., both preferentially act on fatty acids esterified to the sn-2 position of phosphatidylcholine, both use cytochrome b5 as an intermediate electron donor, both are inhibited by cyanide, both require molecular oxygen as a substrate, both are thought to be located in the endoplasmic reticulum).

The third argument stems from the discussion of oxygenase

cofactors above, in which it is suggested that the plant membrane bound fatty acid desaturases may have a μ -oxo bridged diiron cluster-type cofactor, and that such cofactors are capable of catalyzing both fatty acid desaturations and hydroxylations, depending upon the electronic and structural properties of the protein active site.

Taking these three arguments together, it was hypothesized that kappa hydroxylase of castor endosperm is homologous to the microsomal oleate Δ 12 desaturase found in all plants. The evidence supporting this hypothesis has been disclosed.

A number of genes encoding microsomal Δ 12 desaturases from various species have recently been cloned (Okuley et al., 1994) and substantial information about the structure of these enzymes is now known. Hence, in the following invention we teach how to use structural information about fatty acyl desaturases to isolate kappa hydroxylase genes of this invention. This example teaches the method by which any carbon-monoxide insensitive plant fatty acyl hydroxylase gene can be identified by one skilled in the art.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-D show the mass spectra of hydroxy fatty acids standards (Figure 1A, O-TMS-methylricinoleate; Figure 1B, O-TMS-methyl densipoleate; Figure 1C, O-TMS-methyl-lesqueroleate; and Figure 1D, O-TMS-methylauricoleate.

Figure 2 shows the fragmentation pattern of

trim thylsilylated methyl esters of hydroxy fatty acids.

Figure 3A shows the gas chromatogram of fatty acids extracted from seeds of wild type *Arabidopsis* plants. Figure 3B shows the gas chromatogram of fatty acids extracted from seeds of transgenic *Arabidopsis* plants containing the *fah12* hydroxylase gene. The numbers indicate the following fatty acids: [1] 16:0; [2] 18:0; [3] 18:1cis Δ 9; [4] 18:2^{cis Δ 9,12}; [5] 20:0; [6] 20:1^{cis Δ 11}; [7] 18:3^{cis Δ 9,12,15}; [8] 22:1^{cis Δ 13}; [9] 24:1^{cis Δ 13}; [10] ricinoleic acid; [11] densipolic acid; [12] lesquerolic acid; [13] auricolic acid.

Figures 4A-D show the mass spectra of novel fatty acids found in seeds of transgenic plants. Figure 4A shows the mass spectrum of peak 10 from Figure 3B. Figure 4B shows the mass spectrum of peak 11 from Figure 3B. Figure 4C shows the mass spectrum of peak 12 from Figure 3B. Figure 4D shows the mass spectrum of peak 13 from Figure 3B.

Figure 5 shows the nucleotide sequence of pLesq2 (SEQ ID NO:1).

Figure 6 shows the nucleotide sequence of pLesq3 (SEQ ID NO:2).

Figure 7 shows a Northern blot of total RNA from seeds of *L. fendleri* probed with pLesq2 or pLesq3. S, indicates RNA is from seeds; L, indicates RNA is from leaves.

Figures 8A-B show the nucleotide sequence of genomic clone encoding pLesq-HYD (SEQ ID NO:3), and the deduced amino acid sequence of hydroxylase enzyme encoded by the gene (SEQ ID NO:4).

Figures 9A-B show multiple sequence alignment of deduced

amino acid sequences for kappa hydroxylases and microsomal $\Delta 12$ desaturases. Abbreviations are: Rcfah12, fah12 hydroxylase gene from *R. communis* (van de Loo et al., 1995); Lffah12, kappa hydroxylase gene from *L. fendleri*; Atfad2, fad2 desaturase from *Arabidopsis thaliana* (Okuley et al., 1994); Gmfad2-1, fad2 desaturase from *Glycine max* (GenBank accession number L43920); Gmfad2-2, fad2 desaturase from *Glycine max* (Genbank accession number L43921); Zmfad2, fad2 desaturase from *Zea mays* (WO 94/11516); Rcfad2, fragment of fad2 desaturase from *R. communis* (WO 94/11516); Bnfad2, fad2 desaturase from *Brassica napus* (WO/ 94/11516); LFFAH12.AMI, SEQ ID NO:4; FAH12.AMI, SEQ ID NO:5; ATFAD2.AMI, SEQ ID NO:6; BNFAD2.AMI, SEQ ID NO:7; GMFAD2-1.AMI, SEQ ID NO:8; GMFAD2-2.AMI, SEQ ID NO:9; ZMFAD2.AMI, SEQ ID NO:10; and RCFAD2.AMI, SEQ ID NO:11.

Figure 10 shows a Southern blot of genomic DNA from *L. fendleri* probed with pLesq-HYD. E=EcoRI, H = HindIII, X = XbaI.

Figure 11 shows a map of binary Ti plasmid pSLJ44024.

SUMMARY OF THE INVENTION

This invention relates to plant fatty acyl hydroxylases. Methods to use conserved amino acid or nucleotide sequences to obtain plant fatty acyl hydroxylases are described. Also described is the use of cDNA clones encoding a plant hydroxylase to produce a family of hydroxylated fatty acids in

transgenic plants.

In a first embodiment, this invention is directed to recombinant DNA constructs which can provide for the transcription or transcription and translation (expression) of the plant kappa hydroxylase sequence. In particular, constructs which are capable of transcription or transcription and translation in plant host cells are preferred. Such constructs may contain a variety of regulatory regions including transcriptional initiation regions obtained from genes preferentially expressed in plant seed tissue. In a second aspect, this invention relates to the presence of such constructs in host cells, especially plant host cells which have an expressed plant kappa hydroxylase therein.

In yet another aspect, this invention relates to a method for producing a plant kappa hydroxylase in a host cell or progeny thereof via the expression of a construct in the cell. Cells containing a plant kappa hydroxylase as a result of the production of the plant kappa hydroxylase encoding sequence are also contemplated herein.

In another embodiment, this invention relates to methods of using a DNA sequence encoding a plant kappa hydroxylase for the modification of the proportion of hydroxylated fatty acids produced within a cell, especially plant cells. Plant cells having such a modified hydroxylated fatty acid composition are also contemplated herein.

In a further aspect of this invention, plant kappa hydroxylase proteins and sequences which are related thereto, including amino acid and nucleic acid sequences, are

contemplated. Plant kappa hydroxylase exemplified herein includes a *Lesquerella fendleri* fatty acid hydroxylase. This exemplified fatty acid hydroxylase may be used to obtain other plant fatty acid hydroxylases of this invention.

5 In a further aspect of this invention, a nucleic acid sequence which directs the seed specific expression of an associated polypeptide coding sequence is described. The use of this nucleic acid sequence or fragments derived thereof, to obtain seed-specific expression in higher plants of any coding
10 sequence is contemplated herein.

DETAILED DESCRIPTION OF THE INVENTION

A genetically transformed plant of the present invention
15 which accumulates hydroxylated fatty acids can be obtained by expressing the double-stranded DNA molecules described in this application.

A plant fatty acid hydroxylase of this invention includes any sequence of amino acids, such as a protein, polypeptide or
20 peptide fragment, or nucleic acid sequences encoding such polypeptides, obtainable from a plant source which demonstrates the ability to catalyze the production of ricinoleic, lesquerolic, hydroxyerucic (16-hydroxydocos-cis-13-enoic acid) or hydroxypalmitoleic (12-hydroxyhexadec-cis-9-
25 enoic) from CoA, ACP or lipid-linked monoenoic fatty acid substrates under plant enzyme reactive conditions. By "enzyme reactive conditions" is meant that any necessary conditions are available in an environment (i.e., such factors as

temperature, pH, lack of inhibiting substances) which will permit the enzyme to function.

Preferential activity of a plant fatty acid hydroxylase toward a particular fatty acyl substrate is determined upon comparison of hydroxylated fatty acid product amounts obtained per different fatty acyl substrates. For example, by "oleate preferring" is meant that the hydroxylase activity of the enzyme preparation demonstrates a preference for oleate-containing substrates over other substrates. Although the precise substrate of the castor fatty acid hydroxylase is not known, it is thought to be a monounsaturated fatty acid moiety which is esterified to a phospholipid such as phosphatidylcholine. However, it is also possible that monounsaturated fatty acids esterified to phosphatidylethanolamine, phosphatidic acid or a neutral lipid such as diacylglycerol or a Coenzyme-A thioester may also be substrates. As noted above, significant activity has been observed in radioactive labelling studies using fatty acyl substrates other than oleate (Howling et al., 1972) indicating that the substrate specificity is for a family of related fatty acyl compounds. Because the castor hydroxylase introduces hydroxy groups three carbons from a double bond, proximal to the methyl carbon of the fatty acid we term the enzyme a kappa hydroxylase for convenience. Of particular interest, we envision that the castor kappa hydroxylase may be used for production of 12-hydroxy-9-octadecenoic acid (ricinoleate), 12-hydroxy-9-hexadecenoic acid, 14-hydroxy-11-eicosenoic acid, 16-hydroxy-13-docosanoic acid, 9-hydroxy-6-

octadecenoic acid by expression in plants species which produce the non-hydroxylated precursors. We also envision production of additionally modified fatty acids such as 12-hydroxy-9,15-octadecadienoic acid that result from
5 desaturation of hydroxylated fatty acids (e.g., 12-hydroxy-9-octadecenoic acid in this example).

We also envision that future advances in the genetic engineering of plants will lead to production of substrate fatty acids, such as icosenoic acid esters, and palmitoleic
10 acid esters in plants that do not normally accumulate such fatty acids. We envision that the invention described herein may be used in conjunction with such future improvements to produce hydroxylated fatty acids of this invention in any plant species that is amenable to directed genetic
15 modification. Thus, the applicability of this invention is not limited in our conception only to those species that currently accumulate suitable substrates.

As noted above, a plant kappa hydroxylase of this invention will display activity towards various fatty acyl
20 substrates. During biosynthesis of lipids in a plant cell, fatty acids are typically covalently bound to acyl carrier protein (ACP), coenzyme A (CoA) or various cellular lipids. Plant kappa hydroxylases which display preferential activity toward lipid-linked acyl substrate are especially preferred
25 because they are likely to be closely associated with normal pathway of storage lipid synthesis in immature embryos. However, activity toward acyl-CoA substrates or other synthetic substrates, for example, is also contemplated

herein.

Other plant kappa hydroxylases are obtainable from the specific exemplified sequences provided herein. Furthermore, it will be apparent that one can obtain natural and synthetic plant kappa hydroxylases including modified amino acid sequences and starting materials for synthetic-protein modeling from the exemplified plant kappa hydroxylase and from plant kappa hydroxylases which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences which have been mutated, truncated, increased and the like, whether such sequences were partially or wholly synthesized. Sequences which are actually purified from plant preparations or are identical or encode identical proteins thereto, regardless of the method used to obtain the protein or sequence, are equally considered naturally derived.

Thus, one skilled in the art will readily recognize that antibody preparations, nucleic acid probes (DNA and RNA) and the like may be prepared and used to screen and recover "homologous" or "related" kappa hydroxylases from a variety of plant sources. Typically, nucleic acid probes are labeled to allow detection, preferably with radioactivity although enzymes or other methods may also be used. For immunological screening methods, antibody preparations either monoclonal or polyclonal are utilized. Polyclonal antibodies, although less specific, typically are more useful in gene isolation. For detection, the antibody is labeled using radioactivity or any one of a variety of second antibody/enzyme conjugate systems that are commercially available.

Homologous sequences are found when there is an identity of sequence and may be determined upon comparison of sequence information, nucleic acid or amino acid, or through hybridization reactions between a known kappa hydroxylase and a candidate source. Conservative changes, such as Glu/Asp, Val/Ile, Ser/Thr, Arg/Lys and Gln/Asn may also be considered in determining sequence homology. Typically, a lengthy nucleic acid sequence may show as little as 50-60% sequence identity, and more preferably at least about 70% sequence identity, between the target sequence and the given plant kappa hydroxylase of interest excluding any deletions which may be present, and still be considered related. Amino acid sequences are considered homologous by as little as 25% sequence identity between the two complete mature proteins. (See generally, Doolittle, R.F., OF URFS and ORFS, University Science Books, CA, 1986.)

A genomic or other appropriate library prepared from the candidate plant source of interest may be probed with conserved sequences from the plant kappa hydroxylase to identify homologically related sequences. Use of an entire cDNA or other sequence may be employed if shorter probe sequences are not identified. Positive clones are then analyzed by restriction enzyme digestion and/or sequencing. When a genomic library is used, one or more sequences may be identified providing both the coding region, as well as the transcriptional regulatory elements of the kappa hydroxylase gene from such plant source. Probes can also be considerably shorter than the entire sequence. Oligonucleotides may be

used, for example, but should be at least about 10, preferably at least about 15, more preferably at least 20 nucleotides in length. When shorter length regions are used for comparison, a higher degree of sequence identity is required than for longer sequences. Shorter probes are often particularly useful for polymerase chain reactions (PCR), especially when highly conserved sequences can be identified (See Gould, et al., 1989 for examples of the use of PCR to isolate homologous genes from taxonomically diverse species).

When longer nucleic acid fragments are employed (>100 bp) as probes, especially when using complete or large cDNA sequences, one would screen with low stringencies (for example, 40-50°C below the melting temperature of the probe) in order to obtain signal from the target sample with 20-50% deviation, i.e., homologous sequences. (Beltz, et al. 1983).

In a preferred embodiment, a plant kappa hydroxylase of this invention will have at least 60% overall amino acid sequence similarity with the exemplified plant kappa hydroxylase. In particular, kappa hydroxylases which are obtainable from an amino acid or nucleic acid sequence of a castor or lesquerella kappa hydroxylase are especially preferred. The plant kappa hydroxylases may have preferential activity toward longer or shorter chain fatty acyl substrates. Plant fatty acyl hydroxylases having oleate-12-hydroxylase activity and eicosenoate-14-hydroxylase activity are both considered homologously related proteins because of in vitro evidence (Howling et al., 1972), and evidence disclosed herein, that the castor kappa hydroxylase will act on both

substrates. Hydroxylated fatty acids may be subject to further enzymatic modification by other enzymes which are normally present or are introduced by genetic engineering methods. For example, 14-hydroxy-11,17-eicosadienoic acid, which is present in some *Lesquerella* species (Smith 1985), is thought to be produced by desaturation of 14-hydroxy-11-eicosenoic acid.

Again, not only can gene clones and materials derived thereof be used to identify homologous plant fatty acyl hydroxylases, but the resulting sequences obtained therefrom may also provide a further method to obtain plant fatty acyl hydroxylases from other plant sources. In particular, PCR may be a useful technique to obtain related plant fatty acyl hydroxylases from sequence data provided herein. One skilled in the art will be able to design oligonucleotide probes based upon sequence comparisons or regions of typically highly conserved sequence. Of special interest are polymerase chain reaction primers based on the conserved regions of amino acid sequence between the castor kappa hydroxylase and the *L. fendleri* hydroxylase (SED ID NO:4). Details relating to the design and methods for a PCR reaction using these probes are described more fully in the examples.

It should also be noted that the fatty acyl hydroxylases of a variety of sources can be used to investigate fatty acid hydroxylation events in a wide variety of plant and in vivo applications. Because all plants synthesize fatty acids via a common metabolic pathway, the study and/or application of one plant fatty acid hydroxylase to a heterologous plant host may

be readily achieved in a variety of species.

Once the nucleic acid sequence is obtained, the transcription, or transcription and translation (expression), of the plant fatty acyl hydroxylases in a host cell is desired to produce a ready source of the enzyme and/or modify the composition of fatty acids found therein in the form of free fatty acids, esters (particularly esterified to glycerolipids or as components of wax esters), estolides, or ethers. Other useful applications may be found when the host cell is a plant host cell, in vitro and in vivo.

For example, by increasing the amount of an kappa hydroxylase available to the plant, an increased percentage of ricinoleate or lesqueroleate (14-hydroxy-11-eicosenoic acid) may be provided.

Kappa hydroxylase

By this invention, a mechanism for the biosynthesis of ricinoleic acid in plants is demonstrated. Namely, that a specific plant kappa hydroxylase having preferential activity toward fatty acyl substrates is involved in the accumulation of hydroxylated fatty acids in at least some plant species. The use of the terms ricinoleate or ricinoleic acid (or lesqueroleate or lesquerolic acid, densipoleate etc.) is intended to include the free acids, the ACP and CoA esters, the salts of these acids, the glycerolipid esters (particularly the triacylglycerol esters), the wax esters, the estolides and the ether derivatives of these acids.

The determination that plant fatty acyl hydroxylases are active in the *in vivo* production of hydroxylated fatty acids suggests several possibilities for plant enzyme sources. And in fact, hydroxylated fatty acids are found in some natural plant species in abundance. For example, three hydroxy fatty acids related to ricinoleate occur in major amounts in seed oils from various *Lesquerella* species. Of particular interest, lesquerolic acid is a 20 carbon homolog of ricinoleate with two additional carbons at the carboxyl end of the chain (Smith 1985). Other natural plant sources of hydroxylated fatty acids include but are not limited to seeds of the *Linum* genus, seeds of *Wrightia* species, *Lycopodium* species, *Strophanthus* species, *Convolvulaceae* species, *Calendula* species and many others (van de Loo et al., 1993).

Plants having significant presence of ricinoleate or lesqueroleate or desaturated other or modified derivatives of these fatty acids are preferred candidates to obtain naturally-derived kappa hydroxylases. For example, *Lesquerella densipila* contains a diunsaturated 18 carbon fatty acid with a hydroxyl group (van de Loo et al., 1993) that is thought to be produced by an enzyme that is closely related to the castor kappa hydroxylase, according to the theory on which this invention is based. In addition, a comparison between kappa hydroxylases and between plant fatty acyl hydroxylases which introduce hydroxyl groups at positions other than the 12-carbon of oleate or the 14-carbon of lesqueroleate or on substrates other than oleic acid and icosanoic acid may yield insights for protein modeling or other modifications to create

synthetic hydroxylases as discussed above. For example, on the basis of information gained from structural comparisons of the $\Delta 12$ desaturases and the kappa hydroxylase, we envision making genetic modifications in the structural genes for $\Delta 12$ desaturases that convert these desaturases to kappa-hydroxylases. We also envision making changes in $\Delta 15$ hydroxylases that convert these to hydroxylases with comparable substrate specificity to the desaturases (e.g., conversion of $18:2^{\Delta 9,12}$ to $15OH-18:2^{\Delta 9,12}$. Since the difference between a hydroxylase and a desaturases concerns the disposition of one proton, we envision that by systematically changing the charged groups in the region of the enzyme near the active site, we can effect this change.

Especially of interest are fatty acyl hydroxylases which demonstrate activity toward fatty acyl substrates other than oleate, or which introduce the hydroxyl group at a location other than the C12 carbon. As described above, other plant sources may also provide sources for these enzymes through the use of protein purification, nucleic acid probes, antibody preparations, protein modeling, or sequence comparisons, for example, and of special interest are the respective amino acid and nucleic acid sequences corresponding to such plant fatty acyl hydroxylases. Also as previously described, once a nucleic acid sequence is obtained for the given plant hydroxylase, further plant sequences may be compared and/or probed to obtain homologously related DNA sequences thereto and so on.

Genetic Engineering Applications

As is well known in the art, once a cDNA clone encoding a plant kappa hydroxylase is obtained, it may be used to obtain
5 its corresponding genomic nucleic acid sequences thereto.

The nucleic acid sequences which encode plant kappa hydroxylases may be used in various constructs, for example, as probes to obtain further sequences from the same or other species. Alternatively, these sequences may be used in
10 conjunction with appropriate regulatory sequences to increase levels of the respective hydroxylase of interest in a host cell for the production of hydroxylated fatty acids or study of the enzyme *in vitro* or *in vivo* or to decrease or increase levels of the respective hydroxylase of interest for some
15 applications when the host cell is a plant entity, including plant cells, plant parts (including but not limited to seeds, cuttings or tissues) and plants.

A nucleic acid sequence encoding a plant kappa hydroxylase of this invention may include genomic, cDNA or
20 mRNA sequence. By "encoding" is meant that the sequence corresponds to a particular amino acid sequence either in a sense or anti-sense orientation. By "recombinant" is meant that the sequence contains a genetically engineered modification through manipulation via mutagenesis, restriction
25 enzymes, and the like. A cDNA sequence may or may not encode pre-processing sequences, such as transit or signal peptide sequences. Transit or signal peptide sequences facilitate the delivery of the protein to a given organelle and are

frequently cleaved from the polypeptide upon entry into the organelle, releasing the "mature" sequence. The use of the precursor DNA sequence is preferred in plant cell expression cassettes.

5 Furthermore, as discussed above the complete genomic sequence of the plant kappa hydroxylase may be obtained by the screening of a genomic library with a probe, such as a cDNA probe, and isolating those sequences which regulate expression in seed tissue. In this manner, the transcription and
10 translation initiation regions, introns, and/or transcript termination regions of the plant kappa hydroxylase may be obtained for use in a variety of DNA constructs, with or without the kappa hydroxylase structural gene. Thus, nucleic acid sequences corresponding to the plant kappa hydroxylase of
15 this invention may also provide signal sequences useful to direct transport into an organelle 5' upstream non-coding regulatory regions (promoters) having useful tissue and timing profiles, 3' downstream non-coding regulatory region useful as transcriptional and translational regulatory regions and may
20 lend insight into other features of the gene.

 Once the desired plant kappa hydroxylase nucleic acid sequence is obtained, it may be manipulated in a variety of ways. Where the sequence involves non-coding flanking regions, the flanking regions may be subjected to resection,
25 mutagenesis, etc. Thus, transitions, transversions, deletions, and insertions may be performed on the naturally occurring sequence. In addition, all or part of the sequence may be synthesized. In the structural gene, one or more

codons may be modified to provide for a modified amino acid sequence, or one or more codon mutations may be introduced to provide for a convenient restriction site or other purpose involved with construction or expression. The structural gene
5 may be further modified by employing synthetic adapters, linkers to introduce one or more convenient restriction sites, or the like.

The nucleic acid or amino acid sequences encoding a plant kappa hydroxylase of this invention may be combined with other
10 non-native, or "heterologous", sequences in a variety of ways. By "heterologous" sequences is meant any sequence which is not naturally found joined to the plant kappa hydroxylase, including, for example, combination of nucleic acid sequences from the same plant which are not naturally found joined
15 together.

The DNA sequence encoding a plant kappa hydroxylase of this invention may be employed in conjunction with all or part of the gene sequences normally associated with the kappa
hydroxylase. In its component parts, a DNA sequence encoding
20 kappa hydroxylase is combined in a DNA construct having, in the 5' to 3' direction of transcription, a transcription initiation control region capable of promoting transcription and translation in a host cell, the DNA sequence encoding plant kappa hydroxylase and a transcription and translation
25 termination region.

Potential host cells include both prokaryotic and eukaryotic cells. A host cell may be unicellular or found in a multicellular differentiated or undifferentiated organism

depending upon the intended use. Cells of this invention may be distinguished by having a plant kappa hydroxylase foreign to the wild-type cell present therein, for example, by having a recombinant nucleic acid construct encoding a plant kappa hydroxylase therein.

Depending upon the host, the regulatory regions will vary, including regions from viral, plasmid or chromosomal genes, or the like. For expression in prokaryotic or eukaryotic microorganisms, particularly unicellular hosts, a wide variety of constitutive or regulatable promoters may be employed. Expression in a microorganism can provide a ready source of the plant enzyme. Among transcriptional initiation regions which have been described are regions from bacterial and yeast hosts, such as *E. coli*, *B. subtilis*, *Saccharomyces cerevisiae*, including genes such as beta-galactosidase, T7 polymerase, tryptophan E and the like.

For the most part, the constructs will involve regulatory regions functional in plants which provide for modified production of plant kappa hydroxylase with resulting modification of the fatty acid composition. The open reading frame, coding for the plant kappa hydroxylase or functional fragment thereof will be joined at its 5' end to a transcription initiation regulatory region such as the wild-type sequence naturally found 5' upstream to the kappa hydroxylase structural gene. Numerous other transcription initiation regions are available which provide for a wide variety of constitutive or regulatable, e.g., inducible, transcription of the structural gene functions. Among

transcriptional initiation regions used for plants are such regions associated with the structural genes such as for nopaline and mannopine synthases, or with napin, soybean β -conglycinin, oleosin, 12S storage protein, the cauliflower mosaic virus 35S promoters and the like. The transcription/translation initiation regions corresponding to such structural genes are found immediately 5' upstream to the respective start codons. In embodiments wherein the expression of the kappa hydroxylase protein is desired in a plant host, the use of all or part of the complete plant kappa hydroxylase gene is desired; namely all or part of the 5' upstream non-coding regions (promoter) together with the structural gene sequence and 3' downstream non-coding regions may be employed. If a different promoter is desired, such as a promoter native to the plant host of interest or a modified promoter, i.e., having transcription initiation regions derived from one gene source and translation initiation regions derived from a different gene source, including the sequence encoding the plant kappa hydroxylase of interest, or enhanced promoters, such as double 35S CaMV promoters, the sequences may be joined together using standard techniques.

For such applications when 5' upstream non-coding regions are obtained from other genes regulated during seed maturation, those preferentially expressed in plant embryo tissue, such as transcription initiation control regions from the *B. napus* napin gene, or the Arabidopsis 12S storage protein, or soybean β -conglycinin (Bray et al., 1987), or the *L. fendleri* kappa hydroxylase promoter described herein are

desired. Transcription initiation regions which are preferentially expressed in seed tissue, i.e., which are undetectable in other plant parts, are considered desirable for fatty acid modifications in order to minimize any disruptive or adverse effects of the gene product.

Regulatory transcript termination regions may be provided in DNA constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding the plant kappa hydroxylase or a convenient transcription termination region derived from a different gene source, for example, the transcript termination region which is naturally associated with the transcript initiation region. Where the transcript termination region is from a different gene source, it will contain at least about 0.5 kb, preferably about 1-3 kb of sequence 3' to the structural gene from which the termination region is derived.

Plant expression or transcription constructs having a plant kappa hydroxylase as the DNA sequence of interest for increased or decreased expression thereof may be employed with a wide variety of plant life, particularly, plant life involved in the production of vegetable oils for edible and industrial uses. Most especially preferred are temperate oilseed crops. Plants of interest include, but are not limited to rapeseed (Canola and high erucic acid varieties), Crambe, *Brassica juncea*, *Brassica nigra*, meadowfoam, flax, sunflower, safflower, cotton, *Cuphea*, soybean, peanut, coconut and oil palms and corn. An important criterion in the selection of suitable plants for the introduction on the kappa

hydroxylase is the presence in the host plant of a suitable substrate for the hydroxylase. Thus, for example, production of ricinoleic acid will be best accomplished in plants that normally have high levels of oleic acid in seed lipids.

5 Similarly, production of lesquerolic acid will best be accomplished in plants that have high levels of icosanoic acid in seed lipids.

Depending on the method for introducing the recombinant constructs into the host cell, other DNA sequences may be
10 required. Importantly, this invention is applicable to dicotyledons and monocotyledons species alike and will be readily applicable to new and/or improved transformation and regulation techniques. The method of transformation is not
15 critical to the current invention; various methods of plant transformation are currently available. As newer methods are available to transform crops, they may be directly applied hereunder. For example, many plant species naturally susceptible to *Agrobacterium* infection may be successfully transformed via tripartite or binary vector methods of
20 *Agrobacterium* mediated transformation. In addition, techniques of microinjection, DNA particle bombardment, electroporation have been developed which allow for the transformation of various monocot and dicot plant species.

In developing the DNA construct, the various components
25 of the construct or fragments thereof will normally be inserted into a convenient cloning vector which is capable of replication in a bacterial host, e.g., *E. coli*. Numerous vectors exist that have been described in the literature.

After each cloning, the plasmid may be isolated and subjected to further manipulation, such as restriction, insertion of new fragments, ligation, deletion, insertion, resection, etc., so as to tailor the components of the desired sequence. Once the
5 construct has been completed, it may then be transferred to an appropriate vector for further manipulation in accordance with the manner of transformation of the host cell.

Normally, included with the DNA construct will be a structural gene having the necessary regulatory regions for
10 expression in a host and providing for selection of transformant cells. The gene may provide for resistance to a cytotoxic agent, e.g., antibiotic, heavy metal, toxin, etc., complementation providing prototrophy to an auxotrophic host, viral immunity or the like. Depending upon the number of
15 different host species the expression construct or components thereof are introduced, one or more markers may be employed, where different conditions for selection are used for the different hosts.

It is noted that the degeneracy of the DNA code provides
20 that some codon substitutions are permissible of DNA sequences without any corresponding modification of the amino acid sequence.

As mentioned above, the manner in which the DNA construct is introduced into the plant host is not critical to this
25 invention. Any method which provides for efficient transformation may be employed. Various methods for plant cell transformation include the use of Ti- or Ri-plasmids, microinjection, electroporation, infiltration, imbibition, DNA

particle bombardment, liposome fusion, DNA bombardment or the like. In many instances, it will be desirable to have the construct bordered on one or both sides of the T-DNA, particularly having the left and right borders, more particularly the right border. This is particularly useful when the construct uses *A. tumefaciens* or *A. rhizogenes* as a mode for transformation, although the T-DNA borders may find use with other modes of transformation.

Where *Agrobacterium* is used for plant cell transformation, a vector may be used which may be introduced into the *Agrobacterium* host for homologous recombination with T-DNA or the Ti- or Ri-plasmid present in the *Agrobacterium* host. The Ti- or Ri-plasmid containing the T-DNA for recombination may be armed (capable of causing gall formation) or disarmed (incapable of causing gall), the latter being permissible, so long as the *vir* genes are present in the transformed *Agrobacterium* host. The armed plasmid can give a mixture of normal plant cells and gall.

In some instances where *Agrobacterium* is used as the vehicle for transforming plant cells, the expression construct bordered by the T-DNA border(s) will be inserted into a broad host spectrum vector, there being broad host spectrum vectors described in the literature. Commonly used is pRK2 or derivatives thereof. See, for example, Ditta et al., (1980).

Included with the expression construct and the T-DNA will be one or more markers, which allow for selection of transformed *Agrobacterium* and transformed plant cells. A number of

markers have been developed for use with plant cells, such as resistance to kanamycin, the aminoglycoside G418, hygromycin, or the like. The particular marker employed is not essential to this invention, one or another marker being preferred depending on the particular host and the manner of construction.

For transformation of plant cells using *Agrobacterium*, explants may be combined and incubated with the transformed *Agrobacterium* for sufficient time for transformation, the bacteria killed, and the plant cells cultured in an appropriate selective medium. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be grown to seed and the seed used to establish repetitive generations and for isolation of vegetable oils.

The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

EXAMPLES

In the experimental disclosure which follows, all temperatures are given in degrees centigrade ($^{\circ}$), weights are given in grams (g), milligram (mg) or micrograms (μ g), concentrations are given as molar (M), millimolar (mM) or micromolar (μ M) and all volumes are given in liters (l),

microliters (μ l) or milliliters (ml), unless otherwise indicated.

5 EXAMPLE 1 - PRODUCTION OF NOVEL HYDROXYLATED FATTY ACIDS IN
 ARABIDOPSIS THALIANA

Overview

 The kappa hydroxylase encoded by the previously described
 fah12 gene from Castor (U.S. Patent application 08/320,982)
10 was used to produce ricinoleic acid, lesquerolic acid,
 densipolic acid and auricolic acid in transgenic Arabidopsis
 plants. This example reduces to practice the method taught in
 Example 2 of the foregoing application.

15 Production of transgenic plants

 A variety of methods have been developed to insert a DNA
 sequence of interest into the genome of a plant host to obtain
 the transcription and translation of the sequence to eff ct
 phenotypic changes. The following methods represent only one
20 of many equivalent means of producing transgenic plants and
 causing expression of the hydroxylase gene.

 Arabidopsis plants were transformed, by *Agrobacterium*-
 mediated transformation, with the kappa hydroxylase encoded by
 the Castor fah12 gene on binary Ti plasmid pB6. This plasmid
25 was previously used to transform *Nicotiana tabacum* for the
 production of ricinoleic acid (U.S. Patent application
 08/320,982).

 Inoculums of *Agrobacterium tumefaciens* strain GV3101

containing binary Ti plasmid pB6 were plated on L-broth plates containing 50 µg/ml kanamycin and incubated for 2 days at 30°C. Single colonies were used to inoculate large liquid cultures (L-broth medium with 50 mg/l rifampicin, 110 mg/l gentamycin and 200 mg/l kanamycin) to be used for the transformation of Arabidopsis plants.

Arabidopsis plants were transformed by the *in planta* transformation procedure essentially as described by Bechtold et al., (1993). Cells of *A. tumefaciens* GV3101(pB6) were harvested from liquid cultures by centrifugation, then resuspended in infiltration medium at OD₆₀₀ = 0.8 (Infiltration medium was Murashige and Skoog macro and micronutrient medium (Sigma Chemical Co., St. Louis, MO) containing 10 mg/l 6-benzylaminopurine and 5% glucose). Batches of 12-15 plants were grown for 3 to 4 weeks in natural light at a mean daily temperature of approximately 25°C in 3.5 inch pots containing soil. The intact plants were immersed in the bacterial suspension then transferred to a vacuum chamber and placed under 600 mm of vacuum produced by a laboratory vacuum pump until tissues appeared uniformly water-soaked (approximately 10 min). The plants were grown at 25°C under continuous light (100 µmol m⁻² s⁻¹ irradiation in the 400 to 700 nm range) for four weeks. The seeds obtained from all the plants in a pot were harvested as one batch. The seeds were sterilized by sequential treatment for 2 min with ethanol followed by 10 min in a mixture of household bleach (Chlorox), water and Tween-80 (50%, 50%, 0.05%) then rinsed thoroughly with sterile water. The seeds were plated at high density (2000 to 4000 per plate)

onto agar-solidified medium in 100 mm petri plates containing 1/2 X Murashige and Skoog salts medium enriched with B5 vitamins (Sigma Chemical Co., St. Louis, MO) and containing kanamycin at 50 mg/l. After incubation for 48 h at 4°C to stimulate germination, seedlings were grown for a period of seven days until transformants were clearly identifiable as healthy green seedlings against a background of chlorotic kanamycin-sensitive seedlings. The transformants were transferred to soil for two weeks before leaf tissue could be used for DNA and lipid analysis. More than 20 transformants were obtained.

DNA was extracted from young leaves from transformants to verify the presence of an intact fah12 gene. The presence of the transgene in a number of the putative transgenic lines was verified by using the polymerase chain reaction to amplify the insert from pB6. The primers used were HF2 = GCTCTTTTGTGCGCTCATTC (SEQ ID NO:12) and HR1 = CGGTACCAGAAAACGCCTTG (SEQ ID NO:13), which were designed to allow the amplification of a 700 bp fragment. Approximately 100 ng of genomic DNA was added to a solution containing 25 pmol of each primer, 1.5 U Taq polymerase (Boehringer Mannheim), 200 uM of dNTPs, 50 mM KCl, 10 mM Tris.Cl (pH 9), 0.1% (v/v) Triton X-100, 1.5 mM MgCl₂, 3% (v/v) formamide, to a final volume of 50 µl. Amplifications conditions were: 4 min denaturation step at 94°C, followed by 30 cycles of 92°C for 1 min, 55°C for 1 min, 72°C for 2 min. A final extension step closed the program at 72°C for 5 min. Transformants could be positively identified after visualization of a characteristic

1 kb amplified fragment on an ethidium bromide stained agarose gel. All transgenic lines tested gave a PCR product of a size consistent with the expected genotype, confirming that the lines were, indeed, transgenic. All further experiments were done with three representative transgenic lines of the wild type designated as 1-3, 4D, 7-4 and one transgenic line of the fad2 mutant line JB12. The transgenic JB12 line was included in order to test whether the increased accumulation of oleic acid in this mutant would have an effect on the amount of ricinoleic acid that accumulated in the transgenic plants.

Analysis of transgenic plants

Leaves and seeds from fah12 transgenic Arabidopsis plants were analyzed for the presence of hydroxylated fatty acids using gas chromatography. Lipids were extracted from 100-200 mg leaf tissue or 50 seeds. Fatty acid methyl esters (FAMES) were prepared by placing tissue in 1.5 ml of 1.0 M methanolic HCl (Supelco Co.) in a 13 x 100 mm glass screw-cap tube capped with a teflon-lined cap and heated to 80°C for 2 hours. Upon cooling, 1 ml petroleum ether was added and the FAMES removed by aspirating off the ether phase which was then dried under a nitrogen stream in a glass tube. One hundred μ l of N,O-bis(trimethylsilyl) trifluoroacetamide (BSTFA; Pierce Chemical Co) and 200 μ l acetonitrile was added to derivatize the hydroxyl groups. The reaction was carried out at 70°C for 15 min. The products were dried under nitrogen, redissolved in 100 μ l chloroform and transferred to a gas chromatograph vial. Two μ l of each sample were analyzed on a SP2340 fused

silica capillary column (30 m, 0.75 mm ID, 0.20 mm film, Supelco), using a Hewlett-Packard 5890 II series Gas Chromatograph. The samples were not split, the temperature program was 195°C for 18 min, increased to 230°C at 25°C/min, held at 230°C for 5 min then down to 195°C at 25°C/min., and flame ionization detectors were used.

The chromatographic elution time of methyl esters and O-TMS derivatives of ricinoleic acid, lesquerolic acid and auricolic acid was established by GC-MS of lipid samples from seeds of *L. fendleri* and comparison to published chromatograms of fatty acids from this species (Carlson et al., 1990). A O-TMS-methyl-ricinoleate standard was prepared from ricinoleic acid obtained from Sigma Chemical Co (St, Louis, MO). O-TMS-methyl-lesqueroleate and O-TMS-methyl-auricoleate standards were prepared from triacylglycerols purified from seeds of *L. fendleri*. The mass spectrum of O-TMS-methyl-ricinoleate, O-TMS-methyl-densipoleate, O-TMS-methyl-lesqueroleate, and O-TMS-methyl-auricoleate are shown in Figures 1A-D, respectively. The structures of the characteristic ions produced during mass spectrometry of these derivatives are shown in Figure 2.

Lipid extracted from transgenic tissues were analyzed by gas chromatography and mass spectrometry for the presence of hydroxylated fatty acids. As a matter of reference, the average fatty acid composition of leaves in *Arabidopsis* wild type and *fad2* mutant lines was reported by Miquel and Browse (1992). Gas chromatograms of methylated and silylated fatty acids from seeds of wild type and a *fah12* transgenic wild type

plant are shown in Figures 3A and 3B, respectively. The profiles are very similar except for the presence of three small but distinct peaks at 14.3, 15.9 and 18.9 minutes. A very small peak at 20.15 min was also evident. The elution time of the peaks at 14.3 and 18.9 min corresponded precisely to that of comparably prepared ricinoleic and lesquerolic standards, respectively. No significant differences were observed in lipid extracts from leaves or roots of the wild type and the fah12 transgenic wild type lines (Table 1). Thus, in spite of the fact that the fah12 gene is expressed throughout the plant, we observed effects on fatty acid composition only in seed tissue. A similar observation was described previously for transgenic fah12 tobacco in patent application No. 08/320,982.

Table 1: Fatty acid composition of lipids from transgenic and wild type *Arabidopsis*. The values are the means obtained from analysis of samples from three independent transgenic lines, or three independent samples of wild type and *fad2* lines.

Fatty acid	Seed				Leaf		Root	
	WT	FAH12/WT	FAH12/ <i>fad2</i>	JB12	WT	FAH12/WT	WT	FAH12/WT
16:0	8.5	8.2	6.4	6.1	16.5	17.5	23.9	24.9
16:3	0	0	0	0	10.1	9.8	0	0
18:0	3.2	3.5	2.9	3.5	1.3	1.2	2.0	1.9
18:1	15.4	26.3	43.4	47.8	2.4	3.4	5.4	3.2
18:2	27.0	21.4	10.2	7.2	15.1	14.0	32.2	29.4
18:3	22.0	16.6	-	9.7	36.7	36.0	26.7	30.6
20:1	14.0	14.3	-	13.1	0	0	0	0
22:1	2.0	1.0	0.5	0.5	0	0	0	0
24:1	2.5	1.7	2.0	1.6	0	0	0	0
18:1-OH	0	0.4	0.3	0	0	0	0	0
18:2-OH	0	0.4	0.3	0	0	0	0	0
20:1-OH	0	0.2	0.1	0	0	0	0	0
20:2-OH	0	0.1	0.1	0	0	0	0	0

In order to confirm that the observed new peaks in the transgenic lines corresponded to derivatives of ricinoleic, lesquerolic, densipolic and auricolic acids, mass spectrometry was used. The fatty acid derivatives were resolved by gas chromatography as described above except that a Hewlett-Packard 5971 series mass selective detector was used in place of the flame ionization detector used in the previous experiment. The spectra of the four new peaks in Figure 3B (peak numbers 10, 11, 12 and 13) are shown in Figures 4A-D, respectively. Comparison of the spectrum obtained for the standards with that obtained for the four peaks from the transgenic lines confirms the identity of the four new peaks. On the basis of the three characteristic peaks at M/Z 187, 270 and 299, peak 10 is unambiguously identified as O-TMS-methylricinoleate. On the basis of the three characteristic peaks at M/Z 185, 270 and 299, peak 11 is unambiguously identified as O-TMS-methyldensipoleate. On the basis of the three characteristic peaks at M/Z 187, 298 and 327, peak 12 is unambiguously identified as O-TMS-methyllesqueroleate. On the basis of the three characteristic peaks at M/Z 185, 298 and 327, peak 13 is unambiguously identified as O-TS-methylauricoleate. These results unequivocally demonstrate the identity of the fah12 cDNA as encoding a hydroxylase that hydroxylates both oleic acid to produce ricinoleic acid and also hydroxylate icosenoic acid to produce lesquerolic acid. These results also provide additional evidence that the hydroxylase can be functionally expressed in a heterologous plant species in such a way that the enzyme is catalytically

functional. These results also demonstrate that expression of this hydroxylase gene leads to accumulation of ricinoleic, lesquerolic, densipolic and auricolic acids in a plant species that does not normally accumulate hydroxylated fatty acids in extractable lipids.

The presence of lesquerolic acid in the transgenic plants was anticipated in the previous patent application (No. 08/320,982) based on the biochemical evidence suggesting broad substrate specificity of the kappa hydroxylase. By contrast, the accumulation of densipolic and auricolic acids was less predictable. Since Arabidopsis does not normally contain significant quantities of the non-hydroxylated precursors of these fatty acids which could serve as substrates for the hydroxylase, it appears that one or more of the three n-3 fatty acid desaturases known in Arabidopsis (eg., fad3, fad7, fad8; reviewed in Gibson et al., 1995) are capable of desaturating the hydroxylated compounds at the n-3 position. That is, densipolic acid is produced by the action of an n-3 desaturase on ricinoleic acid. Auricolic acid is produced by the action of an n-3 desaturase on lesquerolic acid. Because it is located in the endoplasmic reticulum, the fad3 desaturase is almost certainly responsible. This can be tested in the future by producing fah12-containing transgenic plants of the fad3-deficient mutant of Arabidopsis (similar experiments can be done with fad7 and fad8). It is also formally possible that the enzymes that normally elongate 18:1^{cisΔ9} to 20:1^{cisΔ11} may elongate 12OH-18:1^{cisΔ9} to 14OH-20:1^{cisΔ11}, and 12OH-18:2^{cisΔ9,15} to 14OH-20:2^{cisΔ11,17}.

The amount of the various fatty acids in seed, leaf and root lipids of the control and transgenic plants is presented in Table 1. Although the amount of hydroxylated fatty acids produced in this example is less than desired for commercial production of ricinoleate and other hydroxylated fatty acids from plants, we envision numerous improvements of this invention that will increase the level of accumulation of hydroxylated fatty acids in plants that express the *fah12* or related hydroxylase genes. Improvements in the level and tissue specificity of expression of the hydroxylase gene is envisioned. Methods to accomplish this by the use of strong, seed-specific promoters such as the *B. napus* napin promoter or the native promoters of the castor *fah12* gene or the corresponding hydroxylase gene from *L. fendleri* will be obvious to one skilled in the art. Additional improvements are envisioned to involve modification of the enzymes which cleave hydroxylated fatty acids from phosphatidylcholine, reduction in the activities of enzymes which degrade hydroxylated fatty acids and replacement of acyltransferases which transfer hydroxylated fatty acids to the sn-1, sn-2 and sn-3 positions of glycerolipids. Although genes for these enzymes have not been described in the scientific literature, their utility in improving the level of production of hydroxylated fatty acids can be readily envisioned based on the results of biochemical investigations of ricinoleate synthesis.

Although *Arabidopsis* is not an economically important plant species, it is widely accepted by plant biologists as a

model for higher plants. Therefore, the inclusion of this example is intended to demonstrate the general utility of the invention described here

to the modification of oil composition in higher plants. One advantage of studying the expression of this novel gene in *Arabidopsis* is the existence in this system of a large body of knowledge on lipid metabolism, as well as the availability of a collection of mutants which can be used to provide useful information on the biochemistry of fatty acid hydroxylation in plant species. Another advantage is the ease of transposing any of the information obtained on metabolism of ricinoleate in *Arabidopsis* to closely related species such as the crop plants *Brassica napus*, *Brassica juncea* or *Crambe Abyssinica* in order to mass produce ricinoleate, lesqueroleate or other hydroxylated fatty acids for industrial use. The kappa hydroxylase is useful for the production of ricinoleate or lesqueroleate in any plant species that accumulates significant levels of the precursors, oleic acid and icosanoic acid. Of particular interest are genetically modified varieties that accumulate high levels of oleic acid. Such varieties are currently available for sunflower and Canola. Production of lesquerolic acid and related hydroxy fatty acids can be achieved in species that accumulate high levels of icosanoic acid or other long chain monoenoic acids. Such plants may in the future be produced by genetic engineering of plants that do not normally make such precursors. Thus, we envision that the use of the kappa hydroxylase is of general utility.

EXAMPLE 2. ISOLATION OF LESQUERELLA KAPPA HYDROXYLASE GENOMIC CLONE

Overview

5 Regions of nucleotide sequence that were conserved in both the
Castor kappa hydroxylase and the Arabidopsis fad2 $\Delta 12$ fatty
acid desaturase were used to design oligonucleotide primers.
These were used with genomic DNA from *Lesquerella fendleri* to
amplify fragments of several homologous genes. These
10 amplified fragments were then used as hybridization probes to
identify full length genomic clones from a genomic library of
L. fendleri. Hydroxylated fatty acids are specific to the
seed tissue of *Lesquerella* sp., and are not found to any
appreciable extent in vegetative tissues. One of the two
15 genes identified by this method was expressed in both leaves
and developing seeds and is therefore thought to correspond to
the $\Delta 12$ fatty acid desaturase. The other gene was expressed
at high levels in developing seeds but was not expressed or
was expressed at very low levels in leaves and is the kappa
20 hydroxylase from this species. The identity of this gene will
be established by introducing the gene into transgenic
Arabidopsis plants and showing that it causes the accumulation
of ricinoleic acid, lesquerolic acid, densipolic acid and
auricolic acid in seed lipids. The promoter of this gene is
25 also of utility because it is able to direct expression of a
gene specifically in developing seeds at a time when storage
lipids are accumulating. This promoter is, therefore, of
great utility for many applications in the genetic engineering

of seeds, particularly in members of the Brassicacea.

The various steps involved in this process are described in detail below. Unless otherwise indicated, routine methods for manipulating nucleic acids, bacteria and phage were as described by Sambrook et al. (1989).

Isolation of a fragment of the Lesquerella kappa hydroxylase gene

Oligonucleotide primers for the amplification of the *L. fendleri* kappa hydroxylase were designed by choosing regions of high deduced amino acid sequence homology between the Castor kappa hydroxylase and the Arabidopsis $\Delta 12$ desaturase (fad2). Because most amino acids are encoded several different codons, these oligonucleotides were designed to encode all possible codons that could encode the corresponding amino acids. The sequence of these mixed oligonucleotides was:

Oligo 1: TAYWSNCA YMGNMGNCAYCA (SEQ ID NO:14)

Oligo 2: RTGRTGNGCNACRTGNGTRTC (SEQ ID NO:15)

(Where: Y = C+T; W = A+T; S = G+C; N = A+G+C+T; M = A+C; R = A+G)

These oligonucleotides were used to amplify a fragment of DNA from *L. fendleri* genomic DNA by the polymerase chain reaction (PCR) using the following conditions: Approximately 100 ng of genomic DNA was added to a solution containing 25 pmol of each primer, 1.5 U Taq polymerase (Boehringer Mannheim), 200 μ M of dNTPs, 50 mM KCl, 10 mM Tris.Cl (pH 9), 0.1% (v/v) Triton X-100, 1.5 mM MgCl₂, 3% (v/v) formamide, to a final volume of 50 μ l. Amplifications conditions were: 4 min

denaturation step at 94°C, followed by 30 cycles of 92°C for 1 min, 55°C for 1 min, 72°C for 2 min. A final extension step closed the program at 72°C for 5 min.

5 PCR products of approximately 540 bp were observed following electrophoretic separation of the products of the PCR reaction in agarose gels. Two of these fragments were cloned into pBluescript (Stratagene) to give rise to plasmids pLesq2 and pLesq3. The sequence of the inserts in these two plasmids was determined by the chain termination method. The
10 sequence of the insert in pLesq2 is presented as Figure 5 (SEQ ID NO:1) and the sequence of the insert in pLesq3 is presented as Figure 6 (SEQ ID NO:2). The high degree of sequence identity between the two clones indicated that they were both potential candidates to be either a $\Delta 12$ desaturase or a gamma
15 hydroxylase.

Northern analysis

In *L. fendleri*, hydroxylated fatty acids are found in large amounts in seed oils but are not found in appreciable
20 amounts in leaves. Therefore, an important criterion in discriminating between a fatty acyl desaturase and kappa hydroxylase is that the kappa hydroxylase gene is expected to be expressed more highly in tissues which have high level of hydroxylated fatty acids than in other tissues whereas all
25 plant tissues should contain mRNA for an $\omega 6$ fatty acyl desaturase since diunsaturated fatty acids are found in the lipids of all tissues in most or all plants. Therefore, it was of great interest to determine whether the gene

corresponding to pLesq2 was also expressed only in seeds, or is also expressed in other tissues. This question was addressed by testing for hybridization of pLesq2 to RNA purified from developing seeds and from leaves.

5 Total RNA was purified from developing seeds and young leaves of *L. fendleri* using an Rneasy RNA extraction kit (Qiagen), according to the manufacturer's instructions. RNA concentrations were quantified by UV spectrophotometry at $\lambda=260$ and 280 nm. In order to ensure even loading of the gel
10 to be used for Northern blotting, RNA concentrations were further adjusted after recording fluorescence under UV light of RNA samples stained with ethidium bromide and run on a test denaturing gel.

 Total RNA prepared as described above from leaves and
15 developing seeds was electrophoresed through an agarose gel containing formaldehyde (Iba et al., 1993). An equal quantity (10 μ g) of RNA was loaded in both lanes, and RNA standards (0.16-1.77 kb ladder, Gibco-BRL) were loaded in a third lane. Following electrophoresis, RNA was transferred from the gel to
20 a nylon membrane (Hybond N+, Amersham) and fixed to the filter by exposure to UV light. A 32 P-labelled probe was prepared from insert DNA of clone pLesq2 by random priming and hybridized to the membrane overnight at 52°C, after it had been prehybridized for 2 h. The prehybridization solution
25 contained 5X SSC, 10X Denhardt's solution, 0.1% SDS, 0.1M KPO₄ pH 6.8, 100 μ g/ml salmon sperm DNA. The hybridization solution had the same basic composition, but no SDS, and it contained 10% dextran sulfate and 30% formamide. The blot was

washed once in 2X SSC, 0.5% SDS at 65°C then in 1X SSC at the same temperature.

Brief (30 min) exposure of the blot to X-ray film revealed that the probe pLesq2 hybridized to a single band only in the seed RNA lane (Figure 7). The blot was re-probed with the insert from pLesq3 gene, which gave bands of similar intensity in the seed and leaf lanes (Figure 7).

These results show that the gene corresponding to the clone pLesq2 is highly and specifically expressed in seed of *L. fendleri*. In conjunction with knowledge of the nucleotide and deduced amino acid sequence, strong seed-specific expression of the gene corresponding to the insert in pLesq2 is a convincing indicator of the role of the enzyme in synthesis of hydroxylated fatty acids in the seed oil.

Characterization of a genomic clone of the gamma hydroxylase

Genomic DNA was prepared from young leaves of *L. fendleri* as described by Murray and Thompson (1980). A *Sau3AI*-partial digest genomic library constructed in the vector λ DashII (Stratagene, 11011 North Torrey Pines Road, La Jolla CA 92037) was prepared by partially digesting 500 μ g of DNA, size-selecting the DNA on a sucrose gradient (Sambrook et al., 1989), and ligating the DNA (12 kb average size) to the *Bam*HI-digested arms of λ DashII. The entire ligation was packaged according to the manufacturer's conditions and plated on *E. coli* strain XL1-Blue MRA-P2 (Stratagene). This yielded 5×10^5 primary recombinant clones. The library was then amplified according to the manufacturer's conditions. A fraction of the

g n mic library was plated on *E. coli* XL1-Blue and resulting
plaques (150,000) were lifted to charged nylon membranes
(Hybond N⁺, Amersham), according to the manufacturer's
conditions. DNA was crosslinked to the filters under UV in a
5 Stratalinker (Stratagene).

Several clones carrying genomic sequences corresponding
to the *L. fendleri* hydroxylase were isolated by probing the
membranes with the insert from pLesq2 that was PCR-amplified
with internal primers and labelled with ³²P by random priming.
10 The filters were prehybridized for 2 hours at 65°C in 7% SDS,
1mM EDTA, 0.25 M Na₂HPO₄ (pH 7.2), 1% BSA and hybridized to
the probe for 16 hours in the same solution. The filters were
sequentially washed at 65°C in solutions containing 2 X SSC, 1
X SSC, 0.5 X SSC in addition to 0.1 % SDS. A 2.6 kb *Xba* I
15 fragment containing the complete coding sequence for the
gamma-hydroxylase and approximately 1 kb of the 5' upstream
region was subcloned into the corresponding site of
pBluescript KS to produce plasmid pLesq-Hyd and the sequence
determined completely using an automatic sequencer by the
20 dideoxy chain termination method. Sequence data was analyzed
using the program DNASIS (Hitachi company).

The sequence of the insert in clone pLesq-Hyd is shown in
Figures 8A-B. The sequence entails 1855 bp of contiguous DNA
sequence (SEQ ID NO:3). The clone encodes a 401 bp 5'
25 untranslated region (i.e., nucleotides preceding the first ATG
codon), an 1152 bp open reading frame, and a 302 bp 3'
untranslated region. The open reading frame encodes a 384
amino acid protein with a predicted molecular weight of 44,370

(SEQ ID NO:4). The amino terminus lacks features of a typical signal peptide (von Heijne, 1985).

The exact translation-initiation methionine has not been experimentally determined, but on the basis of deduced amino acid sequence homology to the Castor kappa hydroxylase (n ted below) is thought to be the methionine encoded by the first ATG codon at nucleotide 402.

Comparison of the pLesq-Hyd deduced amino acid sequence with sequences of membrane-bound desaturases and the castor hydroxylase (Figures 9A-B) indicates that pLesq-Hyd is homologous to these genes. This figure shows an alignment of the *L. fendleri* hydroxylase (SEQ ID NO:4) with the castor hydroxylase (van de Loo et al. 1995), the Arabidopsis *fad2* cDNA which encodes an endoplasmic reticulum-localized $\Delta 12$ desaturase (called *fad2*) (Okuley et al., 1994), two soybean *fad2* desaturase clones, a *Brassica napus* *fad2* clone, a *Z a mays* *fad2* clone and partial sequence of a *R. communis* *fad2* clone. The high degree of sequence homology indicates that the gene products are of similar function. For instance, the overall homology between the Lesquerella hydroxylase and the Arabidopsis *fad2* desaturase was 92.2% similarity and 84.8% identity and the two sequences differed in length by only one amino acid.

Southern hybridization

Southern analysis was used to examine the copy number of the genes in the *L. fendleri* genome corresponding to the clone

pLesq-Hyd. Genomic DNA (5 μ g) was digested with *EcoR* I, *Hind* III and *Xba* I and separated on a 0.9% agarose gel. DNA was alkali-blotted to a charged nylon membrane (Hybond N+, Amersham), according to the manufacturer's protocol. The blot was prehybridized for 2 hours at 65°C in 7% SDS, 1mM EDTA, 0.25 M Na₂HPO₄ (pH 7.2), 1% BSA and hybridized to the probe for 16 hours in the same solution with pLesq-Hyd insert PCR-amplified with internal primers and labelled with ³²P by random priming. The filters were sequentially washed at 65°C in solutions containing 2 X SSC, 1 X SSC, 0.5 X SSC in addition to 0.1 % SDS, then exposed to X-ray film.

The probe hybridized with a single band in each digest of *L. fendleri* DNA (Figure 10), indicating that the gene from which pLesq-Hyd was transcribed is present in a single copy in the *L. fendleri* genome.

Expression of pLesq-Hyd in Transgenic Plants

There are a wide variety of plant promoter sequences which may be used to cause tissue-specific expression of cloned genes in transgenic plants. For instance, the napin promoter and the acyl carrier protein promoters have previously been used in the modification of seed oil composition by expression of an antisense form of a desaturase (Knutson et al. 1992). Similarly, the promoter for the β -subunit of soybean β -conglycinin has been shown to be highly active and to result in tissue-specific expression in transgenic plants of species other than soybean (Bray et al., 1987). Thus, although we describe the use of the *L. fendleri*

kappa hydroxylase promoter in the examples described here, other promoters which lead to seed-specific expression may also be employed for the production of modified seed oil composition. Such modifications of the invention described here will be obvious to one skilled in the art.

Constructs for expression of *L. fendleri* kappa hydroxylase in plant cells are prepared as follows: A 13 kb *SalI* fragment containing the pLesq-Hyg gene was ligated into the *XhoI* site of binary Ti plasmid vector pSLJ44026 (Jones et al., 1992) (Figure 11) to produce plasmid pTi-Hyd and transformed into *Agrobacterium tumefaciens* strains GV3101 by electroporation. Strain GV3101 (Koncz and Schell, 1986) contains a disarmed Ti plasmid. Cells for electroporation were prepared as follows. GV3101 was grown in LB medium with reduced NaCl (5 g/l). A 250 ml culture was grown to $OD_{600} = 0.6$, then centrifuged at 4000 rpm (Sorvall GS-A rotor) for 15 min. The supernatant was aspirated immediately from the loose pellet, which was gently resuspended in 500 ml ice-cold water. The cells were centrifuged as before, resuspended in 30 ml ice-cold water, transferred to a 30 ml tube and centrifuged at 5000 rpm (Sorvall SS-34 rotor) for 5 min. This was repeated three times, resuspending the cells consecutively in 30 ml ice-cold water, 30 ml ice-cold 10% glycerol, and finally in 0.75 ml ice-cold 10% glycerol. These cells were aliquoted, frozen in liquid nitrogen, and stored at -80C. Electroporations employed a Biorad Gene pulsar instrument using cold 2 mm-gap cuvettes containing 40 μ l cells and 1 μ l of DNA in water, at a voltage of 2.5 KV, and 200 Ohms

resistance. The electroporated cells were diluted with 1 ml SOC medium (Sambrook et al., 1989, page A2) and incubated at 28°C for 2-4 h before plating on medium containing kanamycin (50 mg/l).

5 *Arabidopsis thaliana* can be transformed with the *Agrobacterium* cells containing pTi-Hyd as described in Example 1 above. Similarly, the presence of hydroxylated fatty acids in the transgenic *Arabidopsis* plants can be demonstrated by the methods described in Example 1 above.

0 Constitutive expression of the *L. fendleri* hydroxylase in transgenic plants

A 1.5 kb EcoR I fragment from pLesq-Hyg comprising the entire coding region of the hydroxylase was gel purified, then
5 cloned into the corresponding site of pBluescript KS (Stratagene). Plasmid DNA from a number of recombinant clones was then restricted with Pst I, which should cut only once in the insert and once in the vector polylinker sequence. Release of a 920 bp fragment with Pst I indicated the right
10 orientation of the insert for further manipulations. DNA from one such clone was further restricted with SalI, the 5' overhangs filled-in with the Klenow fragment of DNA polymerase I, then cut with Sac I. The insert fragment was gel purified, and cloned between the Sma I and Sac I sites of pBI121
15 (Clontech) behind the Cauliflower Mosaic Virus 35S promoter. After checking that the sequence of the junction between insert and vector DNA was appropriate, plasmid DNA from a

recombinant clone was used to transform *A. tumefaciens* (GV3101). Kanamycin resistant colonies were then used for *in planta* transformation of *A. thaliana* as previously described.

5 DNA was extracted from kanamycin resistant seedlings and used to PCR-amplify selected fragments from the hydroxylase using nested primers. When fragments of the expected size could be amplified, corresponding plants were grown in the greenhouse or on agar plates, and fatty acids extracted from fully expanded leaves, roots and dry seeds. GC-MS analysis
10 was then performed as previously described to characterize the different fatty acid species and detect accumulation of hydroxy fatty acids in transgenic tissues.

EXAMPLE 3 - OBTAINING OTHER PLANT FATTY ACYL HYDROXYLASES

15 In a previous patent application, we described the ways in which the castor *fah12* sequence could be used to identify other kappa hydroxylases by methods such as PCR and heterologous hybridization. However, because of the high degree of sequence similarity between $\Delta 12$ desaturases and
20 kappa hydroxylases, prior art does not teach how to distinguish between the two kinds of enzymes without a functional test such as demonstrating activity in transgenic plants or another suitable host (e.g., transgenic microbial or animal cells). The identification of the *L. fendleri*
25 hydroxylase provided for the development of criteria by which a hydroxylase and a desaturase may be distinguished solely on the basis of deduced amino acid sequence information.

Figures 9A-B show a sequence alignment of the castor and

L. fendleri hydroxylase sequences with the castor hydroxylase sequence and all publically available sequences for all plant microsomal $\Delta 12$ fatty acid desaturases. Of the 384 amino acid residues in the castor hydroxylase sequence, more than 95% are identical to the corresponding residue in at least one of the desaturase sequences. Therefore, none of these residues are responsible for the catalytic differences between the hydroxylase and the desaturases. Of the remaining 16 residues in the castor hydroxylase and 14 residues in the *Lesquerella* hydroxylase, all but six represent instances where the hydroxylase sequence has a conservative substitution compared with one or more of the desaturase sequences, or there is wide variability in the amino acid at that position in the various desaturases. By conservative, we mean that the following amino acids are functionally equivalent : Ser/Thr, Ile/ Leu/ Val/ Met, Asp/ Glu. Thus, these structural differences also cannot account for the catalytic differences between the desaturases and hydroxylases. This leaves just six amino acid residues where both the castor hydroxylase and the *Lesquerella* hydroxylase differ from all of the known desaturases and where all of the known microsomal $\Delta 12$ desaturases have the identical amino acid residue. These residues occur at positions 69, 111, 155, 226, 304 and 331 of the alignment in Figure 9. Therefore, these six sites distinguish hydroxylases from desaturases. Based on this analysis, we claim that any enzyme with greater than 60% sequence identity to one of the enzymes listed in Figure 9 can be classified as a hydroxylase if it differs from the sequence of the desaturases at these six

positions. Because of slight differences in the number of residues in a particular protein, the numbering may vary from protein to protein but the intent of the numbering system will be evident if the protein in question is aligned with the castor hydroxylase using the numbering system shown herein. Thus, in conjunction with the methods for using the lesquerella hydroxylase gene to isolate homologous genes, the structural criterion disclosed here teaches how to isolate and identify plant kappa hydroxylase genes for the purpose of genetically modifying fatty acid composition.

In considering which of the six substitutions are solely or primarily responsible for the difference in catalytic activity of the hydroxylases of this invention and the desaturases, we consider it likely that the substitution of a Phe for a Tyr at position 226 may be solely responsible for this difference in catalytic activity because of the known participation of tyrosine radicals in enzyme catalysis. Other substitutions, such as the Ala for Ser at position 331 may have effects at modulating the overall rate of the reaction. On this basis we envision creating novel kappa hydroxylases by site directed mutagenesis of $\Delta 12$ desaturases. We also envision converting $\Delta 15$ desaturases and $\Delta 9$ desaturases to hydroxylases by similar use of site-directed mutagenesis.

CONCLUDING REMARKS

By the above examples, demonstration of critical factors in the production of novel hydroxylated fatty acids by

expression of a kappa hydroxylase gene from Castor in transgenic plants is described. In addition, a complete cDNA sequence of the *Lesquerella fendleri* kappa hydroxylase is also provided. A full sequence of the castor hydroxylase is also given with various constructs for use in host cells. Through this invention, one can obtain the amino acid and nucleic acid sequences which encode plant fatty acyl hydroxylases from a variety of sources and for a variety of applications.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

REFERENCES:

- 5 Beltz, G.A., Jacobs, K.A., Eickbuch, T.H., Cherbas, P.T.,
Kafatos, F.C. (1983) Isolation of multigene families and
determination of homologies by filter hybridization methods.
Methods in Enzymology 100, 266-285.
- 10 Bray, E.A., Naito, S., Pan, N.S., Anderson, E., Dube, P.,
Beachy, R.N. (1987) Expression of the β -subunit of β -
conglycinin in seeds of transgenic plants. Planta 172:364-370.
- 15 Carlson, K.D., Chaudhry, A., Bagby, M.O. (1990) Analysis of
oil and meal from *lesquerella fendleri* seed. J. Am. Oil Chem.
Soc. 67, 438-442.
- 20 Ditta, G., Stanfield, S., Corbin, D., Helinski, D.R. (1980)
Broad host range DNA cloning system for gram-negative
bacteria: Construction of a gene bank of *Rhizobium meliloti*.
Proc. Natl. Acad. Sci. USA 77, 7347-7351.
- 25 Gould, S.J., Subramani, S., Scheffler, I.E. (1989) Use of the
DNA polymerase chain reaction for homology probing. Proc.
Natl. Acad. Sci. USA 86, 1934-1938.
- Hirsinger, F. (1989) New oil crops, in *Oil Crops of the World*,
Robbelen, G., Downey, K.R., and Ashri, A., Eds., McGraw-Hill,
New York, pp. 518-533.

Howling, D., Morris, L.J., Gurr, M.I., James, A.T. (1972) The specificity of fatty acid desaturases and hydroxylases. The dehydrogenation and hydroxylation of monoenoic acids, *Biochim. Biophys. Acta* 260, 10.

5

Huyuh, T.V., Young, R.A., Davis, R.W. (1985) Constructing and screening cDNA libraries in λ gt10 and λ gt11. In *DNA Cloning*, Vol. 1: A Practical Approach, (ed) D.M. Glover. IRL Press, Washington DC pp 49-77.

3

Iba, K., Gibson, S., Nishiuchi, T., Fuse, T., Nishimura, M., Arondel, V., Hugly, S., and Somerville, C. (1993) A gene encoding a chloroplast omega-3 fatty acid desaturase complements alterations in fatty acid desaturation and chloroplast copy number of the *fad7* mutant of *Arabidopsis thaliana*. *J. Biol. Chem.* 268, 24099-24105.

5

Jones, J.D.G., Shlumukov, L., Carland, F., English, J., Scofield, S., Bishop, G.J., Harrison, K. (1992) Effective vectors for transformation, expression of heterologous genes, and assaying transposon excision in transgenic plants. *Transgenic Res.* 1, 285-297.

)

Knutson, D.S., Thompson, G.A., Radke, S.E., Johnson, W.B., Knauf, V.C., Kridl, J.C. (1992) *Proc. Natl. Acad. Sci. USA* 89, 2624-2628.

;

Koncz, C., Schell, J. (1986) The promoter of T_1 -DNA gene 5

controls the tissue-specific expression of chimeric genes carried by a novel type of *Agrobacterium* binary vector. *Mol. Gen. Genet.* 204, 383-396.

- 5 Miquel, M. Browse, J. (1992) *Arabidopsis* mutants deficient in polyunsaturated fatty acid synthesis. *J. Biol. Chem.* 267, 1502-1509.

- 10 Murray, M.G., Thompson, W.F. (1980) Rapid isolation of high molecular weight plant DNA. *Nucl. Acid Res.* 8,4321-4325.

- Okuley, J., Lightner, J., Feldman, K., Yadav, N., Lark, E., Browse, J. (1994) *Arabidopsis* FAD2 gene encodes the enzyme that is essential for polyunsaturated lipid

15

Sambrook, J., Fritsch, E.F., and Maniatis, T., *Molecular Cloning: a Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, 1989.

20

Smith C.R., Jr. (1985) Unusual seed oils and their fatty acids, in *Fatty Acids*, Pryde E.H., Ed., American Oil Chemists' Society, Champaign, Second edition, pp 29-47.

- 25 van de Loo, F.J., Fox, B.G., Somerville, C. (1993) Unusual fatty acids, in *Lipid Metabolism in Plants*, T.S. Moore Jr., Ed., CRC Press, Boca Raton, pp91-126.

van de Loo, F.N., Turner, S., Somerville, C.R. (1995) An
oleate 12-hydroxylase from castor (*Ricinus communis* L.) is a
fatty acyl desaturase homolog. Proc. Natl. Acad. Sci. USA
92,6743-6747

5 v n Heijne, G. (1985) Signal sequences. J. Mol. Biol. 184,99-
105

Atsmon, D. (1989) Castor, in *Oil Crops of the World*, Robbelen,
G., Downey, K.R., and Ashri, A., Eds., McGraw-Hill, New York,
pp. 438-447.

Bechtold, N., Ellis, J. and Pelletier, G. (1993) In Planta
Agrobacterium mediated gene transfer by infiltration of adult
Arabidopsis thaliana plants. C.R. Acad. Sci. Paris 316,
1194-1199.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Somerville, Chris
Broun, Pierre
- (ii) TITLE OF INVENTION: PRODUCTION OF HYDROXYLATED FATTY ACIDS
IN GENETICALLY MODIFIED PLANTS
- (iii) NUMBER OF SEQUENCES: 15
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: CUSHMAN DARBY & CUSHMAN, LLP
 - (B) STREET: 1100 NEW YORK AVENUE, NW
 - (C) CITY: WASHINGTON
 - (D) STATE: DC
 - (E) COUNTRY: USA
 - (F) ZIP: 20005-3918
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: PCT/US95/11855
 - (B) FILING DATE: 25-SEP-1995
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Kokulis, Paul N.
 - (B) REGISTRATION NUMBER: 16,773
 - (C) REFERENCE/DOCKET NUMBER: 1220/213781
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 202-861-3000
 - (B) TELEFAX: 202-822-0944

(2) INFORMATION FOR SEQ ID NO:1

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 543 nucleotides
 - (B) TYPE: nucleotide
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TATTGGCACC GCGGGCACCA TTCCAACAAT GGATCCCTAG AAAAAGATGA AGTCTTTGTC 60
CCACCTAAGA AAGCTGCAGT CANATGGTAT GTCAAATACC TCAACAACCC TCTTGGACGC 120

ATTCTGGTGT TAACAGTTCA GTTTATCCTC GGGTGGCCTT TGTATCTAGC CTTTAATGTA 180
 TCAGGTAGAC CTTATGATGG TTTGCTTCA CATTCTTCC CTCATGCACC TATCTTTAAG 240
 GACCGTGAAC GTCTCCAGAT ATACATCTCA GATGCTGGTA TTCTAGCTGT CTGTTATGGT 300
 CTTTACCGTT ACGCTGCTTC ACAAGGATTG ACTGCTATGA TCTGCGTCTA CGGAGTACCG 360
 CTTTTGATAG TGAACTTTTT CCTTGTCTTG GTCACCTTCT TGCAGCACAC TCATCCTTCA 420
 TTACCTCACT ATGATTCAAC CGAGTGGGAA TGGATTAGAG GAGCTTTGGT TACGGTAGAC 480
 AGAGACTATG GAATCTTGAA CAAGGTGTTT CACAACATAA CAGACACCCA CGTAGCACAC 540
 CAC 543

(2) INFORMATION FOR SEQ ID NO:2

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 544 nucleotides
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TATAGGCACC GGAGGCACCA TTCCAACACA GGATCCCTCG AAAGAGATGA AGTATTTGTC 60
 CCAAAGCAGA AATCCGCAAT CAACTGGTAC GGCGAATACC TCAACAACCC TCCTGGTCGC 120
 ATCATGATGT TAACTGTCCA GTTCGTCCTC GGATGGCCCT TGTACTTAGC CTTCAACGTT 180
 TCTGGCAGAC CCTACAATGG TTTGCTTCC CATTCTTCC CCAATGCTCC TATCTACAAC 240
 GACCGTGAAC GCCTCCAGAT TTACATCTCT GATGCTGGTA TTCTAGCCGT CTGTTATGGT 300
 CTTTACCGTT ACGCTGTTGC ACAAGGACTA GCCTCAATGA TCTGTCTAAA CGGAGTTCCG 360
 CTTCTGATAG TTAACTTTTT CCTCGTCTTG ATCACTTACT TACAACACAC TCACCCTGCG 420
 TTGCCTCACT ATGATTCATC AGAGTGGGAT TGGCTTAGAG GAGCTTTAGC TACTGTAGAC 480
 AGAGACTATG GAATCTTGAA CAAGGTGTTT CATAACATCA CAGACACCCA CGTCGCACAC 540
 CACT 544

(2) INFORMATION FOR SEQ ID NO:3

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1740 nucleotides
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGAAGCTTT ATAAGAAGTT AGTTTTCTCT GGTGACAGAG AAATTNTGTC AATTGGTAGT	60
GACAGTTGAA GCAACAGGAA CAACAAGGAT GGTGGTGNT GATGCTGATG TGGTGATGTG	120
TTATTCATCA AATACTAAAT ACTACATTAC TTGTTGCTGC CTAATTCTCC TATTTCTCC	180
GCCACCCATT TTGGACCCAC GANCCTTCCA TTAAACCCT CTCTCGTGCT ATTCACCAGA	240
AGAGAAGCCA AGAGAGAGAG AGAGAGAATG TTCTGAGGAT CATTGTCTTC TTCATCGTTA	300
TTAACGTAAG TTTTTTTTGA CCACTCATAT CTAATACTA GTACATGCAA TAGATTAATG	360
ACTGTTCTTT CTTTTGATAT TTTCAGCTTC TTGAATTCAA GATGGGTGCT GGTGGAAGAA	420
TAATGGTTAC CCCCTCTTCC AAGAAATCAG AAAGTGAAGC CCTAAAACGT GGACCATGTG	480
AGAAACCACC ATTCAGTGT AAAGATCTGA AGAAAGCAAT CCCACAGCAT TGTTTCAAGC	540
GCTCTATCCC TCGTTCTTTC TCCTACCTTC TCACAGATAT CACTTTAGTT TCTTGCTTCT	600
ACTACGTTGC CACAAATTAC TTCTCTCTTC TTCCTCAGCC TCTCTCTACT TACCTAGCTT	660
GGCCTCTCTA TTGGGTATGT CAAGGCTGTG TCTTAACCGG TATCTGGGTC ATTGGCCATG	720
AATGTGGTCA CCATGCATTC AGTGACTATC AATGGGTAGA TGACACTGTT GGTTTTATCT	780
TCCATTCTTT CCTTCTCGTC CCTTACTTCT CCTGGAAATA CAGTCATCGT CGTCACCATT	840
CCAACAATGG ATCTCTCGAG AAAGATGAAG TCTTTGTCCC ACCGAAGAAA GCTGCAGTCA	900
AATGGTATGT TAAATACCTC AACAAACCTC TTGGACGCAT TCTGGTGTTA ACAGTTCAGT	960
TTATCCTCGG GTGGCCTTTC TATCTAGCCT TTAATGTATC AGGTAGACCT TATGATGGTT	1020
TCGCTTCACA TTTCTTCCCT CATGCACCTA TCTTTAAAGA CCGAGAACGC CTCCAGATAT	1080
ACATCTCAGA TGCTGGTATT CTAGCTGTCT GTTATGGTCT TTACCGTTAC GCTGCTTCAC	1140
AAGGATTGAC TGCTATGATC TGCGTCTATG GAGTACCGCT TTTGATAGTG AACTTTTTCC	1200
TTGTCTTGGT AACTTTCTTG CAGCACACTC ATCCTTCGTT ACCTCATTAT GATTCAACCG	1260
AGTGGGAATG GATTAGAGGA GCTTTGGTTA CGGTAGACAG AGACTATGGA ATATTGAACA	1320
AGGTGTTCCA TAACATAACA GACACACATG TGGCTCATCA TCTCTTTGCA ACTATACCGC	1380
ATTATAACGC AATGGAAGCT ACAGAGGCGA TAAAGCCAAT ACTTGGTGAT TACTACCACT	1440
TCGATGGAAC ACCGTGGTAT GTGGCCATGT ATAGGGAAGC AAAGGAGTGT CTCTATGTAG	1500
AACCGGATAC GGAACGTGGG AAGAAAGGTG TCTACTATTA CAACAATAAG TTATGAGGCT	1560
GATAGGGCGA GAGAAGTGCA ATTATCAATC TTCATTTCCA TGTTTTAGGT GTCTTGTTA	1620
AGAAGCTATG CTTTGTTTCA ATAATCTCAG AGTCCATNTA GTTGTGTTCT GGTGCATTTT	1680
GCCTAGTTAT GTGGTGTCGG AAGTTAGTGT TCAAAGTGCT TCCTGCTGTG CTGCCAGTG	1740

AAGAACAAGT TTACGTGTTT AAAATACTCG GAACGAATTG ACCACAANAT ATCCAAAACC 1800
 GGCTATCCGA ATTCCATATC CGAAAACCGG ATATCCAAAT TTCCAGAGTA CTTAG 1835

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Gly	Ala	Gly	Gly	Arg	Ile	Met	Val	Thr	Pro	Ser	Ser	Lys	Lys	Ser	1	5	10	15
Glu	Thr	Glu	Ala	Leu	Lys	Arg	Gly	Pro	Cys	Glu	Lys	Pro	Pro	Phe	Thr	20	25	30	
Val	Lys	Asp	Leu	Lys	Lys	Ala	Ile	Pro	Gln	His	Cys	Phe	Lys	Arg	Ser	35	40	45	
Ile	Pro	Arg	Ser	Phe	Ser	Tyr	Leu	Leu	Thr	Asp	Ile	Thr	Leu	Val	Ser	50	55	60	
Cys	Phe	Tyr	Tyr	Val	Ala	Thr	Asn	Tyr	Phe	Ser	Leu	Leu	Pro	Gln	Pro	65	70	75	80
Leu	Ser	Thr	Tyr	Leu	Ala	Trp	Pro	Leu	Tyr	Trp	Val	Cys	Gln	Gly	Cys	85	90	95	
Val	Leu	Thr	Gly	Ile	Trp	Val	Ile	Gly	His	Glu	Cys	Gly	His	His	Ala	100	105	110	
Phe	Ser	Asp	Tyr	Gln	Trp	Val	Asp	Asp	Thr	Val	Gly	Phe	Ile	Phe	His	115	120	125	
Ser	Phe	Leu	Leu	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Tyr	Ser	His	Arg	Arg	130	135	140	
His	His	Ser	Asn	Asn	Gly	Ser	Leu	Glu	Lys	Asp	Glu	Val	Phe	Val	Pro	145	150	155	160
Pro	Lys	Lys	Ala	Ala	Val	Lys	Trp	Tyr	Val	Lys	Tyr	Leu	Asn	Asn	Pro	165	170	175	
Leu	Gly	Arg	Ile	Leu	Val	Leu	Thr	Val	Gln	Phe	Ile	Leu	Gly	Trp	Pro	180	185	190	
Leu	Tyr	Leu	Ala	Phe	Asn	Val	Ser	Gly	Arg	Pro	Tyr	Asp	Gly	Phe	Ala	195	200	205	
Ser	His	Phe	Phe	Pro	His	Ala	Pro	Ile	Phe	Lys	Asp	Arg	Glu	Arg	Leu	210	215	220	

Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu
 225 230 235 240
 Tyr Arg Tyr Ala Ala Ser Gln Gly Leu Thr Ala Met Ile Cys Val Tyr
 245 250 255
 Gly Val Pro Leu Leu Ile Val Asn Phe Phe Leu Val Leu Val Thr Phe
 260 265 270
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Thr Glu Trp
 275 280 285
 Glu Trp Ile Arg Gly Ala Leu Val Thr Val Asp Arg Asp Tyr Gly Ile
 290 295 300
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His
 305 310 315 320
 Leu Phe Ala Thr Ile Pro His Tyr Asn Ala Met Glu Ala Thr Glu Ala
 325 330 335
 Ile Lys Pro Ile Leu Gly Asp Tyr Tyr His Phe Asp Gly Thr Pro Trp
 340 345 350
 Tyr Val Ala Met Tyr Arg Glu Ala Lys Glu Cys Leu Tyr Val Glu Pro
 355 360 365
 Asp Thr Glu Arg Gly Lys Lys Gly Val Tyr Tyr Tyr Asn Asn Lys Leu
 370 375 380

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Thr Ser Asn Asn Ser
 1 5 10 15
 Glu Lys Lys Gly Gly Ser Ser His Leu Lys Arg Ala Pro His Thr Lys
 20 25 30
 Pro Pro Phe Thr Leu Gly Asp Leu Lys Arg Ala Ile Pro Pro His Cys
 35 40 45
 Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Tyr Val Ala Tyr Asp Val
 50 55 60
 Cys Leu Ser Phe Leu Phe Tyr Ser Ile Ala Thr Asn Phe Phe Pro Tyr
 65 70 75 80

Ile Ser Ser Pro Leu Ser Tyr Val Ala Trp Leu Val Tyr Trp Leu Phe
 85 90 95
 Gln Gly Cys Ile Leu Thr Gly Leu Trp Val Ile Gly His Glu Cys Gly
 100 105 110
 His His Ala Phe Ser Glu Tyr Gln Leu Ala Asp Asp Ile Val Gly Leu
 115 120 125
 Ile Val His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser
 130 135 140
 His Arg Arg His His Ser Asn Ile Gly Ser Leu Glu Arg Asp Glu Val
 145 150 155 160
 Phe Val Pro Lys Ser Lys Ser Lys Ile Ser Trp Tyr Ser Lys Tyr Ser
 165 170 175
 Asn Asn Pro Pro Gly Arg Val Leu Thr Leu Ala Ala Thr Leu Leu Leu
 180 185 190
 Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp
 195 200 205
 Arg Phe Ala Cys His Tyr Asp Pro Tyr Gly Pro Ile Phe Ser Glu Arg
 210 215 220
 Glu Arg Leu Gln Ile Tyr Ile Ala Asp Leu Gly Ile Phe Ala Thr Thr
 225 230 235 240
 Phe Val Leu Tyr Gln Ala Thr Met Ala Lys Gly Leu Ala Trp Val Met
 245 250 255
 Arg Ile Tyr Gly Val Pro Leu Leu Ile Val Asn Cys Phe Leu Val Met
 260 265 270
 Ile Thr Tyr Leu Gln His Thr His Pro Ala Ile Pro Arg Tyr Gly Ser
 275 280 285
 Ser Glu Trp Asp Trp Leu Arg Gly Ala Met Val Thr Val Asp Arg Asp
 290 295 300
 Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His Val
 305 310 315 320
 Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu Ala
 325 330 335
 Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp Gly
 340 345 350
 Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu Phe
 355 360 365
 Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr Arg
 370 375 380

Asn Lys Tyr
385

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Gly Ala Gly Gly Arg Met Pro Val Pro Thr Ser Ser Lys Lys Ser
 1             5             10             15

Glu Thr Asp Thr Thr Lys Arg Val Pro Cys Glu Lys Pro Pro Phe Ser
          20             25             30

Val Gly Asp Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser
          35             40             45

Ile Pro Arg Ser Phe Ser Tyr Leu Ile Ser Asp Ile Ile Ile Ala Ser
          50             55             60

Cys Phe Tyr Tyr Val Ala Thr Asn Tyr Phe Ser Leu Leu Pro Gln Pro
          65             70             75             80

Leu Ser Tyr Leu Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val
          85             90             95

Leu Thr Gly Ile Trp Val Ile Ala His Glu Cys Gly His His Ala Phe
          100            105            110

Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser
          115            120            125

Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His
          130            135            140

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys
          145            150            155            160

Gln Lys Ser Ala Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu
          165            170            175

Gly Arg Ile Met Met Leu Thr Val Gln Phe Val Leu Gly Trp Pro Leu
          180            185            190

Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Phe Ala Cys
          195            200            205

His Phe Phe Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln
          210            215            220

Ile Tyr Leu Ser Asp Ala Gly Ile Leu Ala Val Cys Phe Gly Leu Tyr
          225            230            235            240

```

(2) INFORMATION FOR SEQ ID NO:7:

(A) LENGTH: 384 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

Met	Gly	Ala	Gly	Gly	Arg	Met	Gln	Val	Ser	Pro	Pro	Ser	Lys	Lys	Ser
1				5					10					15	
Glu	Thr	Asp	Asn	Ile	Lys	Arg	Val	Pro	Cys	Glu	Thr	Pro	Pro	Phe	Thr
			20					25					30		
Val	Gly	Glu	Leu	Lys	Lys	Ala	Ile	Pro	Pro	His	Cys	Phe	Lys	Arg	Ser
		35					40					45			
Ile	Pro	Arg	Ser	Phe	Ser	His	Leu	Ile	Trp	Asp	Ile	Ile	Ile	Ala	Ser
	50					55					60				
Cys	Phe	Tyr	Tyr	Val	Ala	Thr	Thr	Tyr	Phe	Pro	Leu	Leu	Pro	Asn	Pro
65					70					75					80
Leu	Ser	Tyr	Phe	Ala	Trp	Pro	Leu	Tyr	Trp	Ala	Cys	Gln	Gly	Cys	Val
				85					90					95	
Leu	Thr	Gly	Val	Trp	Val	Ile	Ala	His	Glu	Cys	Gly	His	Ala	Ala	Phe
			100					105					110		

Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser
 115 120 125
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His
 130 135 140
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Arg
 145 150 155 160
 Arg Ser Gln Thr Ser Ser Gly Thr Ala Ser Thr Ser Thr Thr Phe Gly
 165 170 175
 Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr
 180 185 190
 Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala Cys
 195 200 205
 His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln
 210 215 220
 Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu Leu
 225 230 235 240
 Pro Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Leu Arg
 245 250 255
 Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu
 260 265 270
 Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp
 275 280 285
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu
 290 295 300
 Asn Gln Gly Phe His Asn Ile Thr Asp Thr His Glu Ala His His Leu
 305 310 315 320
 Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala Ile
 325 330 335
 Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val Val
 340 345 350
 Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp
 355 360 365
 Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu Xaa
 370 375 380

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 309 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Ser Leu Leu Thr Ser Phe Ser Tyr Val Val Tyr Asp Leu Ser Phe Ala
1           5           10           15

Phe Ile Phe Tyr Ile Ala Thr Thr Tyr Phe His Leu Leu Pro Gln Pro
20           25           30

Phe Ser Leu Ile Ala Trp Pro Ile Tyr Trp Val Leu Gln Gly Cys Leu
35           40           45

Leu Thr Arg Val Cys Gly His His Ala Phe Ser Lys Tyr Gln Trp Val
50           55           60

Asp Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val Pro Tyr
65           70           75           80

Phe Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr Gly Ser
85           90           95

Leu Asp Arg Asp Glu Arg Val Lys Val Ala Trp Phe Ser Lys Tyr Leu
100          105          110

Asn Asn Pro Leu Gly Arg Ala Val Ser Leu Leu Val Thr Leu Thr Ile
115          120          125

Gly Trp Pro Met Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp
130          135          140

Ser Phe Ala Ser His Tyr His Pro Tyr Arg Val Arg Leu Leu Ile Tyr
145          150          155          160

Val Ser Asp Val Ala Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val
165          170          175

Ala Thr Leu Lys Gly Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro
180          185          190

Leu Leu Ile Val Asn Gly Phe Leu Val Thr Ile Thr Tyr Leu Arg Val
195          200          205

His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly Ala Leu Ala Thr
210          215          220

Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His His Ile Thr
225          230          235          240

Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr His
245          250          255

Leu Arg Val Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Asp Thr
260          265          270

Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Arg Glu Cys Leu Tyr Val
275          280          285

```

Glu Pro Asp Glu Gly Thr Ser Glu Lys Gly Val Tyr Trp Tyr Arg Asn
 290 295 300

Lys Tyr Leu Arg Val
 305

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 302 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Phe Ser Tyr Val Val Tyr Asp Leu Thr Ile Ala Phe Cys Leu Tyr Tyr
 1 5 10 15
 Val Ala Thr His Tyr Phe His Leu Leu Pro Gly Pro Leu Ser Phe Arg
 20 25 30
 Gly Met Ala Ile Tyr Trp Ala Val Gln Gly Cys Ile Leu Thr Gly Val
 35 40 45
 Trp Val Val Ala Phe Ser Asp Tyr Gln Leu Leu Asp Asp Ile Val Gly
 50 55 60
 Leu Ile Leu His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr
 65 70 75 80
 Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu
 85 90 95
 Val Phe Val Pro Lys Val Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg
 100 105 110
 Val Leu Thr Leu Ala Val Thr Leu Thr Leu Gly Trp Pro Leu Tyr Leu
 115 120 125
 Ala Leu Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr
 130 135 140
 Asp Pro Tyr Gly Pro Ile Tyr Ser Val Ile Ser Asp Ala Gly Val Leu
 145 150 155 160
 Ala Val Val Tyr Gly Leu Phe Arg Leu Ala Met Ala Lys Gly Leu Ala
 165 170 175
 Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu Val Val Asn Gly Phe
 180 185 190
 Leu Val Leu Ile Thr Phe Leu Gln His Thr His Val Ser Glu Trp Asp
 195 200 205
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu
 210 215 220

Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His Leu
 225 230 235 240
 Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Val Glu Tyr
 245 250 255
 Tyr Arg Phe Asp Glu Thr Pro Phe Val Lys Ala Met Trp Arg Glu Ala
 260 265 270
 Arg Glu Cys Ile Tyr Val Glu Pro Asp Gln Ser Thr Glu Ser Lys Gly
 275 280 285
 Val Phe Trp Tyr Asn Asn Lys Leu Ala Met Glu Ala Thr Val
 290 295 300

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Gly Ala Gly Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu
 1 5 10 15
 Gln Leu Ala Arg Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val
 20 25 30
 Glu Lys Pro Pro Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro
 35 40 45
 His Cys Phe Glu Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His
 50 55 60
 Asp Leu Val Ile Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile
 65 70 75 80
 Pro Ala Leu Pro Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp
 85 90 95
 Ile Ala Gln Gly Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val
 100 105 110
 Gly Leu Val Leu His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys
 115 120 125
 Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp
 130 135 140
 Glu Val Phe Val Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro
 145 150 155 160
 Tyr Val Tyr Asn Asn Pro Val Gly Arg Val Val His Ile Val Val Gln
 165 170 175

Leu Thr Leu Gly Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg
 180 185 190
 Pro Tyr Pro Arg Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr
 195 200 205
 Asn Asp Arg Glu Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val
 210 215 220
 Ala Val Ala Phe Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp
 225 230 235 240
 Trp Val Val Arg Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp
 245 250 255
 Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His
 260 265 270
 Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met
 275 280 285
 Asp Arg Asp Tyr Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp
 290 295 300
 Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala
 305 310 315 320
 Met Glu Ala Thr Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His
 325 330 335
 Phe Asp Pro Thr Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu
 340 345 350
 Cys Ile Tyr Val Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn
 355 360 365
 Lys Lys Phe Xaa
 370

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 224 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln
 1 5 10 15
 Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val
 20 25 30
 Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His His Ser Asn Thr
 35 40 45

Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Lys Ser Ser
 50 55 60
 Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met
 65 70 75 80
 Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe
 85 90 95
 Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro
 100 105 110
 Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser
 115 120 125
 Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile
 130 135 140
 Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu
 145 150 155 160
 Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His
 165 170 175
 Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly
 180 185 190
 Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe
 195 200 205
 His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro
 210 215 220

(2) INFORMATION FOR SEQ ID NO:12

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 nucleotides
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

 (ii) MOLECULE TYPE: DNA

 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GCTCTTTTGT GCGCTCATTC

20

(2) INFORMATION FOR SEQ ID NO:13

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 nucleotides
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CGGTACCAGA AAACGCCTTG

20

(2) INFORMATION FOR SEQ ID NO:14

(i) SEQUENCE CHARACTERISTICS:

(A)	LENGTH:	20 nucleotides
(B)	TYPE:	nucleotide
(C)	STRANDEDNESS:	single
(D)	TOPOLOGY:	linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TAYWSNCAYM GNMGNCA YCA

20

(2) INFORMATION FOR SEQ ID NO:15

(i) SEQUENCE CHARACTERISTICS:

(A)	LENGTH:	21 nucleotides
(B)	TYPE:	nucleotide
(C)	STRANDEDNESS:	single
(D)	TOPOLOGY:	linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

RTGRTGNGCN ACRTGNGTRT C

21

CLAIMS

What is claimed is:

1. An isolated nucleic acid fragment comprising a nucleic acid sequence encoding a fatty acid hydroxylase with an amino acid identity of 60% or greater to the polypeptide encoded by SEQ ID NO:4.

2. The isolated nucleic acid fragment of Claim 1, wherein the amino acid identity is 90% or greater to the polypeptide encoded by SEQ ID NO:4.

3. The isolated nucleic acid fragment of Claim 1, wherein the amino acid identity is 100% of the polypeptide encoded by SEQ ID NO:4.

4. An isolated nucleic acid fragment having a nucleic acid identity of 90% or greater of a nucleotide sequence of SEQ ID NO:1, 2, or 3.

5. An isolated nucleic acid having a nucleotide sequence of SEQ ID NO:1, SEQ ID NO:2 or SEQ ID NO:3.

6. The isolated nucleic acid fragment of Claim 1, wherein said fragment is isolated from an oil-producing plant species.

7. A chimeric gene capable of causing altered levels of ricinoleic acid in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 1, said fragment operably linked to suitable regulatory sequences.

8. A chimeric gene capable of causing altered levels of lesquerolic acid in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 1, said fragment operably linked to suitable regulatory sequences.

9. A chimeric gene capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 1, said fragment operably linked to suitable regulatory sequences.

10. A chimeric gene capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 2, said fragment operably linked to suitable regulatory sequences.

11. A chimeric gene capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 4, said fragment operably linked to suitable regulatory sequences.

12. Plants containing the chimeric gene of any one of claims 7, 8, 9, 10 or 11.

13. Oil obtained from seeds of the plants of claim 12.

14. The isolated nucleic acid fragment of Claim 1, wherein said fragment is obtainable from *Ricinus communis* (L.) (Castor).

15. The isolated nucleic acid fragment of Claim 1, wherein said fragment is obtainable from *Lesquerella fendleri*.

16. A method of producing seed oil containing altered levels of hydroxylated fatty acids comprising:

(a) transforming a plant cell of an oil-producing species with a chimeric gene containing an isolated nucleic acid of Claim 1;

(b) growing fertile plants from the transformed plant cells of step (a);

(c) screening progeny seeds from the fertile plants of step (b) for the desired levels of hydroxylated fatty acids; and

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

17. The method of Claim 16, wherein said crop plant is selected from the group consisting of rapeseed, Crambe, *Brassica juncea*, Canola, flax, sunflower, safflower, cotton, cuphea, soybean, peanut, coconut, oil palm and corn.

18. A method of producing seed oil containing altered levels of hydroxylated fatty acids comprising:

(a) transforming a plant cell of an oil-producing species with a chimeric gene containing the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:2 or SEQ ID NO:3;

(b) growing fertile plants from the transformed plant cells of step (a);

(c) screening progeny seeds from the fertile plants of step (b) for the desired levels of hydroxylated fatty acids; and

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

19. The method of Claim 18, wherein said crop plant is selected from the group consisting of rapeseed, Crambe, *Brassica juncea*, Canola, flax, sunflower, safflower, cotton, cuphea, soybean, peanut, coconut, oil palm and corn.

20. A triglyceride oil from a plant selected from the group consisting of rapeseed, Crambe, *Brassica juncea*, Canola, flax, sunflower, cotton, cuphea, soybean, peanut, coconut, oil palm and corn, wherein the fatty acid composition of the oil has been modified to contain hydroxylated fatty acids by a method comprising growing a plant cell having integrated in its genome a DNA construct containing a plant hydroxylase encoding sequence of Claim 1, under conditions which will permit the transcription and translation of said plant hydroxylase in the plant cells.

21. A method to isolate nucleic acid fragments encoding fatty acid hydroxylases comprising:

(a) comparing SEQ ID NO:4 and other fatty acid hydroxylase sequences and fatty acid desaturases;

(b) identifying conserved sequences of 4 or more amino acids obtained in step (a);


(c) designing degenerate oligomers based on the conserved sequences identified in step (b);

(d) using the degenerate oligomers of step (c) to isolate sequences encoding fatty acid hydroxylases by sequence dependent protocols;


(e) obtaining the deduced amino acid sequence of the encoded gene product from the nucleotide sequence of the gene and;

(f) distinguishing hydroxylase genes from desaturase genes by analyzing amino acid sequence differences between fatty acid desaturases and fatty acid hydroxylases.

Figure 1A




1B




SUBSTITUTE SHEET (RULE 26)

1C




1D




SUBSTITUTE SHEET (RULE 26)


3/15




Ion #2: Mass 299




Ion #3: Mass 270 (characteristic rearrangement ion)



Ion #4: Mass 185 (desaturated analog of Ion #1)



Ion #5: Mass 298 (elongated analog of Ion #3)



Ion #6: Mass 327 (elongated analog of ion





Figure 2

SUBSTITUTE SHEET (RULE 26)



4/15

Figure 3





SUBSTITUTE SHEET (RULE 26)

5/15

Figure 4A Mass spectrum of peak 10 from figure 3B**4B** Mass spectrum of peak 11 from figure 3B

SUBSTITUTE SHEET (RULE 26)

4C **Mass spectrum of peak 12 from figure 3B****4D** **Mass spectrum of peak 13 from figure 3B**

SUBSTITUTE SHEET (RULE 26)

10	20	30	40	50	60
TATTEGCACC	GGCGGCACCA	TTCCAACAAT	GGATCCCTAG	AAAAAGATGA	AGTCTTTGTC
70	80	90	100	110	120
CCACCTAAGA	AAGCTGCAGT	CANATGGTAT	GTCAAATACC	TCAACAACCC	TCTTGGACGC
130	140	150	160	170	180
ATTCTGGTGT	TAACAGTTCA	GTTTATCCTC	GGGTGGCCTT	TGTATCTAGC	CTTTAATGTA
190	200	210	220	230	240
TCAGGTAGAC	CTTATGATGG	TTTCGCTTCA	CATTTCTTCC	CTCATGCACC	TATCTTTAAG
250	260	270	280	290	300
GACCGTGAAC	GTCTCCAGAT	ATACATCTCA	GATGCTGGTA	TTCTAGCTGT	CTGTTATGGT
310	320	330	340	350	360
CTTTACCGTT	ACGCTGCTTC	ACAAGGATTG	ACTGCTATGA	TCTGCGTCTA	CGGAGTACCG
370	380	390	400	410	420
CTTTGATAG	TGAACTTTTT	CCTTGCTTTG	GTCACTTTCT	TGCAGCACAC	TCATCCTTCA
430	440	450	460	470	480
TTACCTCACT	ATGATTCAAC	CGAGTGGGAA	TGGATTAGAG	GAGCTTTGGT	TACGGTAGAC
490	500	510	520	530	540
AGAGCTATG	GAATCTTGAA	CAAGGTGTTT	CACAACATAA	CAGACACCCA	CGTAGCACAC
550					
CAC					

Figure 5

SUBSTITUTE SHEET (RULE 26)

10	20	30	40	50	60
TATAGGCACC	GGAGGCACCA	TTCCAAACACA	GGATCCCTCG	AAAGAGATGA	AGTATTTGTC
70	80	90	100	110	120
CCAAAGCAGA	AATCCGCAAT	CAAGTGCTAC	GGCGAATACC	TCAACAACCC	TCCTGGTCGC
130	140	150	160	170	180
ATCATGATGT	TAAGTGCCA	GTTGGTCTCT	GGATGGCCCT	TGTACTTAGC	CTTCAACGTT
190	200	210	220	230	240
TCTGGCAGAC	CCTACAATGG	TTTCGCTTCC	CATTTCTTCC	CCAATGCTCC	TATCTACAAC
250	260	270	280	290	300
GACCGTGAAC	GCCTCCAGAT	TTACATCTCT	GATGCTGGTA	TTCTAGCCGT	CTGTTATGGT
310	320	330	340	350	360
CTTTACCGTT	ACGCTGTTGC	ACAAGGACTA	GCCTCAATGA	TCTGTCTAAA	CGGAGTTCCG
370	380	390	400	410	420
CTTCTGATAG	TTAACTTTTT	CCTCGTCTTG	ATCACTTACT	TACAACACAC	TCACCCCTGCG
430	440	450	460	470	480
TTGCCTCACT	ATGATTCATC	AGAGTGGGAT	TGGCTTAGAG	GAGCTTTAGC	TACTGTAGAC
490	500	510	520	530	540
AGAGACTATG	GAATCTTGAA	CAAGGTGTTT	CATAACATCA	CAGACACCCA	CGTCGCACAC
550					

CACT

Figure 6

SUBSTITUTE SHEET (RULE 26)

9/15




Figure 7

SUBSTITUTE SHEET (RULE 26)

10/15

AT GAA GGT TTA TAA GAA GTT AGT TTT CTC TGG TGA CAG AGA AAT TNT	47
GTC AAT TGG TAG TGA CAG TTG AAG TAA CAG GAA CAA CAA GGA TGG TGG	95
GTG NTG ATG CTG ATG TGG TGA TGT GTT ATT CAT CAA ATA CTA AAT ACT	143
ACA TTA CTT GTT GGT GGC TAC TCC TCC TAT TTC CTC CGC CAC CCA TTT	191
TGG ACC CAC GAN GGT TCC ATT TAA ACC CTC TCT CGT GGT ATT CAC CAG	239
AAG AGA AGC CAA GAG AGA GAG AGA GAG AAT GTT CTG AGG ATC ATT GTC	287
TTC TTC ATC GTT ATT AAC GTA AGT TTT TTT TGA CCA CTC ATA TCT AAA	335
ATC TAG TAC ATG CAA TAG ATT AAT GAC TGT TCC TTC TTT TGA TAT TTT	383
CAG CTT CTT GAA TTC AAG ATG GGT GCT GGT GGA AGA ATA ATG GTT ACC	431
Pro Ser Ser Lys Lys Ser Glu Thr Glu Ala Leu Lys Arg Gly Pro Cys	26
CCC TCT TCC AAG AAA TCA GAA ACT GAA GCC CTA AAA CGT GGA CCA TGT	479
Glu Lys Pro Pro Phe Thr Val Lys Asp Leu Lys Lys Ala Ile Pro Gln	42
GAG AAA CCA CCA TTC ACT GTT AAA GAT CTG AAG AAA GCA ATC CCA CAG	527
His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe Ser Tyr Leu Leu Thr	58
CAT TGT TTC AAG CGC TCT ATC CCT CGT TCT TTC TCC TAC CTT CTC ACA	575
Asp Ile Thr Leu Val Ser Cys Phe Tyr Tyr Val Ala Thr Asn Tyr Phe	74
GAT ATC ACT TTA GTT TCT TGC TTC TAC TAC GTT GCC ACA AAT TAC TTC	623
Ser Leu Leu Pro Gln Pro Leu Ser Thr Tyr Leu Ala Trp Pro Leu Tyr	90
TCT CTT CTT CCT CAG CCT CTC TCT ACT TAC CTA GGT TGG CCT CTC TAT	671
Trp Val Cys Gln Gly Cys Val Leu Thr Gly Ile Trp Val Ile Gly His	106
TGG GTA TGT CAA GGC TGT GTC TTA ACC GGT ATC TGG GTC ATT GGC CAT	719
Glu Cys Gly His His Ala Phe Ser Asp Tyr Gln Trp Val Asp Asp Thr	122
GAA TGT GGT CAC CAT GCA TTC AGT GAC TAT CAA TGG GTA GAT GAC ACT	767
Val Gly Phe Ile Phe His Ser Phe Leu Leu Val Pro Tyr Phe Ser Trp	138
GTT GGT TTT ATC TTC CAT TCC TTC CTT CTC GTC CCT TAC TTC TCC TGG	815
Lys Tyr Ser His Arg Arg His His Ser Asn Asn Gly Ser Leu Glu Lys	154
AAA TAC AGT CAT CGT CGT CAC CAT TCC AAC AAT GGA TCT CTC GAG AAA	863
Asp Glu Val Phe Val Pro Pro Lys Lys Ala Ala Val Lys Trp Tyr Val	170
GAT GAA GTC TTT GTC CCA CCG AAG AAA GCT GCA GTC AAA TGG TAT GTT	911
Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ile Leu Val Leu Thr Val Gln	186
AAA TAC CTC AAC AAC CCT CTT GGA CGC ATT CTG GTG TTA ACA GTT CAG	959

Figure 8A

SUBSTITUTE SHEET (RULE 26)

11/15

Phe Ile Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg	202
TTT ATC CTC GGG TGG CCT TTG TAT CTA GCC TTT AAT GTA TCA GGT AGA	1007
Pro Tyr Asp Gly Phe Ala Ser His Phe Phe Pro His Ala Pro Ile Phe	218
CCT TAT GAT GGT TTC GCT TCA CAT TTC TTC CCT CAT GCA CCT ATC TTT	1055
Lys Asp Arg Glu Arg Leu Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu	234
AAA GAC CGA GAA CGC CTC CAG ATA TAC ATC TCA GAT GCT GGT ATT CTA	1103
Ala Val Cys Tyr Gly Leu Tyr Arg Tyr Ala Ala Ser Gln Gly Leu Thr	250
GCT GTC TGT TAT GGT CTT TAC CGT TAC GCT GCT TCA CAA GGA TTG ACT	1151
Ala Met Ile Cys Val Tyr Gly Val Pro Leu Leu Ile Val Asn Phe Phe	266
GCT ATG ATC TGC GTC TAT GGA GTA CCG CTT TTG ATA GTG AAC TTT TTC	1199
Leu Val Leu Val Thr Phe Leu Gln His Thr His Pro Ser Leu Pro His	282
CTT GTC TTG GTA ACT TTC TTG CAG CAC ACT CAT CCT TCG TTA CCT CAT	1247
Tyr Asp Ser Thr Glu Trp Glu Trp Ile Arg Gly Ala Leu Val Thr Val	298
TAT GAT TCA ACC GAG TGG GAA TGG ATT AGA GGA GCT TTG GTT ACG GTA	1295
Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp	314
GAC AGA GAC TAT GGA ATA TTG AAC AAG GTG TTC CAT AAC ATA ACA GAC	1343
Thr His Val Ala His His Leu Phe Ala Thr Ile Pro His Tyr Asn Ala	330
ACA CAT GTG GCT CAT CAT CTC TTT GCA ACT ATA CCG CAT TAT AAC GCA	1391
Met Glu Ala Thr Glu Ala Ile Lys Pro Ile Leu Gly Asp Tyr Tyr His	346
ATG GAA GCT ACA GAG GCG ATA AAG CCA ATA CTT GGT GAT TAC TAC CAC	1439
Phe Asp Gly Thr Pro Trp Tyr Val Ala Met Tyr Arg Glu Ala Lys Glu	362
TTC GAT GGA ACA CCG TGG TAT GTG GCC ATG TAT AGG GAA GCA AAG GAG	1487
Cys Leu Tyr Val Glu Pro Asp Thr Glu Arg Gly Lys Lys Gly Val Tyr	378
TGT CTC TAT GTA GAA CCG GAT ACG GAA CGT GGG AAG AAA GGT GTC TAC	1535
Tyr Tyr Asn Asn Lys Leu	384
TAT TAC AAC AAT AAG TTA TGA GGC TGA TAG GGC GAG AGA AGT GCA ATT	1583
ATC AAT CTT CAT TTC CAT GTT TTA GGT GTC TTG TTT AAG AAG CTA TGC	1631
TTT GTT TCA ATA ATC TCA GAG TCC ATN TAG TTG TGT TCT GGT GCA TTT	1679
TGC CTA GTT ATG TGG TGT CGG AAG TTA GTG TTC AAA CTG CTT CCT GCT	1727
GTG CTG CCC AGT GAA GAA CAA GTT TAC GTG TTT AAA ATA CTC GGA ACG	1775
AAT TGA CCA CAA NAT ATC CAA AAC CGG CTA TCC GAA TTC CAT ATC CGA	1823
AAA CCG GAT ATC CAA ATT TCC AGA GTA CTT AG	1855

Figure 8B
SUBSTITUTE SHEET (RULE 26)

12/15

		10	20	30	40	50	
LFFAH12	1	MGAGGRIM--	--VTPSSKKS	--ETEALKRG	PCEKPPFTVK	DLKKAIPQHC	50
FAH12	1	MGGGGRMSTV	ITSNNSEKKG	--GSSHLKRA	PHTKPPFTLG	DLKKAIPPHC	50
ATFAD2	1	MGAGGRMP--	--VPTSSKKS	--ETDITKRV	PCEKPPFSVG	DLKKAIPPHC	50
BNFAD2	1	MGAGGRMO--	--VSPSSKKS	--ETDNKRV	PCETPPFTVG	ELKKAIPPHC	50
GMFAD2-1	1	MGLA-KETTM	GGRGRVAKVE	VQGKKPLSRV	PNTKPPFTVG	QLKKAIPPHC	50
GMFAD2-2	1	MGAGGR----	TQVPPANRKS	--EVDPLKRV	PFEKPOFSL	QIKKAIPPHC	50
ZMFAD2	1	MGAGGRMTEK	EREKOEOLAR	ATGGAAMORS	PVEKPPFTLG	QIKKAIPPHC	50
RCFAD2	1	-----	-----	-----	-----	-----	50
		60	70	80	90	100	
LFFAH12	51	FKRSIPRSFS	YLLTDITLVS	CFYYVATNYF	SLLPOPLSTY	LAWPLYWVCO	100
FAH12	51	FERSFVRSFS	YVAYDVCLSF	LFYSIATNFF	PYISSPLS-Y	VAWLWVWLFQ	100
ATFAD2	51	FKRSIPRSFS	YLISCIIIAS	CFYYVATNYF	SLLPOPLS-Y	LAWPLYWACO	100
BNFAD2	51	FKRSIPRSFS	HLIWDIIIAS	CFYYVATNYF	PLLPNPLS-Y	FAWPLYWACO	100
GMFAD2-1	51	FORSLLTSFS	YVVYDLSFAF	IFY-IATTYF	HLLPOPLS-L	IAPWPIYVWLO	100
GMFAD2-2	51	FORSVLRSFS	YVVYDLTIAF	CLYYVATHYF	HLLPGPLS-F	RGMAIYWAVQ	100
ZMFAD2	51	FERSVLKSFS	YVVHDLVIAA	ALLYFALAII	PALPSPLR-Y	AAWPLYWIAQ	100
RCFAD2	51	-----	-----	-----	-----	-----	100
		110	120	130	140	150	
LFFAH12	101	GCVLTGIWVI	GHECGHHAFS	DYQWVDDTVG	FIFHSFLLVP	YFSWKYSHRR	150
FAH12	101	GCILTGLWVI	GHECGHHAFS	EYQLADDIVG	LIVHSALLVP	YFSWKYSHRR	150
ATFAD2	101	GCVLTGIWVI	AHECGHHAFS	DYQWLDDTVG	LIFHSFLLVP	YFSWKYSHRR	150
BNFAD2	101	GCVLTGVWVI	AHECGHHAFS	DYQWLDDTVG	LIFHSFLLVP	YFSWKYSHRR	150
GMFAD2-1	101	GCLLTGVWVI	AHECGHHAFS	KYQWVDDVVG	LTLHSTLLVP	YFSWKYSHRR	150
GMFAD2-2	101	GCILTGVWVI	AHECGHHAFS	DYQLDDDIVG	LILHSALLVP	YFSWKYSHRR	150
ZMFAD2	101	G-----	-----AFS	OYSLDDVVG	LVLHSSLMVP	YFSWKYSHRR	150
RCFAD2	101	-----WVM	AHDCGHHAFS	DYQLDDVVG	LILHSCLLVP	YFSWKYSHRR	150
		160	170	180	190	200	
LFFAH12	151	HHSNNGSLEK	DEVFVPPKKA	AVKWWVKYL-	NNPLGRILVL	TVOFILGWPL	200
FAH12	151	HHSNIGSLER	DEVFVPPKKS	KISWYSKYS-	NNPPGRVLT	AATLLLGWPL	200
ATFAD2	151	HHSNTGSLER	DEVFVPPKKS	AIKWYGKYL-	NNPLGRIMML	TVOFVLGWPL	200
BNFAD2	151	HHSNTGSLER	DEVFVPR-RS	QTSSGTAST-	STTFGRIVML	TVOFTLGWPL	200
GMFAD2-1	151	HHSNTGSLDR	DEVFVPPKKS	KVAFWSKYL-	NNPLGRAVSL	LVTLTIGWPM	200
GMFAD2-2	151	HHSNTGSLER	DEVFVPPKKS	CIKWYSKYL-	NNPPGRVLT	AVTLTLGWPL	200
ZMFAD2	151	HHSNTGSLER	DEVFVPPKKE	ALPWYTPYVY	NNPVGRVVHI	VVOLTLGWPL	200
RCFAD2	151	HHSNTGSLER	DEVFVPPKKS	SIRWYSKYL-	NNPPGRIMTI	AVTLSLGWPL	200
		210	220	230	240	250	
LFFAH12	201	YLAFNVSGRP	YDG-FASHFF	PHAPIFKDRE	RLOIYISDAG	ILAVCYGLYR	250
FAH12	201	YLAFNVSGRP	YDR-FACHYD	PYGPIFSERE	RLOIYIADLG	IFATTFVLYQ	250
ATFAD2	201	YLAFNVSGRP	YDG-FACHFF	PNAPIYNORE	RLOIYISDAG	ILAVCFGLYR	250
BNFAD2	201	YLAFNVSGRP	YDGGFACHFH	PNAPIYNORE	RLOIYISDAG	ILAVCYGLLP	250
GMFAD2-1	201	YLAFNVSGRP	YDS-FASHYH	PYAPIYSNRE	RLLIYVSDVA	LFSVTYSLYR	250
GMFAD2-2	201	YLALNVSGRP	YDR-FACHYD	PYGPIYSNRE	RLOIYISDAG	VLAVVYGLFR	250
ZMFAD2	201	YLATNASGRP	YPR-FACHFD	PYGPIYNDRE	RAQIFVSDAG	VVAVAFGLYK	250
RCFAD2	201	YLAFNVSGRP	YDR-FACHYD	PYGPIYNORE	RIEIFISDAG	VLAVTFGLYO	250

Figure 9A

SUBSTITUTE SHEET (RULE 26)

		260	270	280	290	300	
LFFAH12	251	AASOGLTAM	ICVYGVPLLI	VNFFLVLVTF	LQHTHPSLPH	YDSTEWEWIR	300
FAH12	251	ATMAKGLAWV	MRIYGVPLLI	VNCFLVMITY	LQHTHPAIPR	YGSSEWDWLR	300
ATFAD2	251	AAAAOQMASH	ICLYGVPLLI	VNAFLVLITY	LQHTHPSLPH	YDSSEWDWLR	300
BNFAD2	251	YAAVQGVASM	VCFLRVPLLI	VNGFLVLITY	LQHTHPSLPH	YDSSEWDWLR	300
GMFAD2-1	251	YATLKGLVWL	LCVYGVPLLI	VNGFLVTITY	LQHTHFALPH	YDSSEWDWLK	300
GMFAD2-2	251	LAMAKGLAWV	VCVYGVPLLV	VNGFLVLITY	LQHTHPALPH	YTSSEWDWLR	300
ZMFAD2	251	LAAGFGVWWV	VRVYAVPLLI	VNAWLVLITY	LQHTHPSLPH	YDSSEWDWLR	300
RCFAD2	251	LAIKGLAWV	VCVYGVPLLV	VNSFLVLITY	LQHTHPALPH	YDSSEWDWLR	300
		310	320	330	340	350	
LFFAH12	301	GALVTVDROY	GILNKVFHNI	TOTHVAHHLF	ATIPHYNAME	ATEAIKPILG	350
FAH12	301	GAMVTVDROY	GVLNKVFHNI	ADTHVAHHLF	ATVPHYHAME	ATKAIKPIMG	350
ATFAD2	301	GALATVDROY	GILNKVFHNI	TOTHVAHHLF	STMPHYNAME	ATKAIKPILG	350
BNFAD2	301	GALATVDROY	GILNOGFHNI	TOTHEAHHLF	STMPHYHAME	ATKAIKPILG	350
GMFAD2-1	301	GALATMDROY	GILNKVFHHI	TOTHVAHHLF	STMPHYHAME	ATNAIKPILG	350
GMFAD2-2	301	GALATVDROY	GILNKVFHNI	TOTHVAHHLF	STMPHYHAME	ATKAIKPILG	350
ZMFAD2	301	GALATMDROY	GILNRVFHNI	TOTHVAHHLF	STMPHYHAME	ATKAIRPILG	350
RCFAD2	301	GALATVDROY	GILNKVFHNI	TOTOVAHHLF	-----	-----	350
		360	370	380	390	400	
LFFAH12	351	DYYHFDGTPW	YVAMYREAKE	CLYVEPDTER	GKKGVYYNN	K-L.....	400
FAH12	351	EYYRYDGTTPF	YKALWREAKE	CLFVEPDEGA	PTQGVFWYRN	KY-.....	400
ATFAD2	351	DYYQFDGTPW	YVAMYREAKE	CIYVEPDREG	DKKGVYWYNN	K-L.....	400
BNFAD2	351	EYYQFDGTPV	VKAMWREAKE	CIYVEPDROG	EKKGVFWYNN	KL*.....	400
GMFAD2-1	351	EYYQFDDTPF	YKALWREARE	CLYVEPDEGT	SEKGVYWYRN	KY-.....	400
GMFAD2-2	351	EYYRFDETPF	VKAMWREARE	CIYVEPDOST	ESKGVFWYNN	KL-.....	400
ZMFAD2	351	DYYHFDPTPV	AKATWREAGE	CIYVEPE---	DRKGVFWYNN	KF*.....	400

Figure 9B

SUBSTITUTE SHEET (RULE 26)

14/15





Figure 10

SUBSTITUTE SHEET (RULE 26)

15/15



Plasmid name: pSLJ44026

Plasmid size: 25.70 kb

Constructed by: Jonathon Jones

Construction date: 1992

Comments/References: Transgenic Research 1,285-297 (1992)

Figure 11

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/11855

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 5/14, 15/29, 15/53, 15/82

US CL : 435/134, 189, 320.1; 800/205

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/134, 189, 320.1; 800/205

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

search terms: fatty acid hydroxylase, oleate hydroxylase, castor.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCIENCE, Volume 258, issued 20 November 1992, Arondel et al., "Map-based cloning of a gene controlling omega-3 fatty acid desaturation in Arabidopsis", pages 1353-1355, see entire document.	1-21
Y	Planta, Volume 172, issued 1987, Bray et al., "Expression of the beta-subunit of beta-conglycinin in seeds of transgenic plants", pages 364-370, see entire document.	1-21
Y	Biochemical Journal, Volume 287, issued 1992, Smith et al., "Evidence for cytochrome b5 as an electron donor in ricinoleic acid biosynthesis in microsomal preparations from developing castor bean (Ricinus communis L.)", pages 141-144, see entire document.	1-21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* Z*	document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means		
* P* documents published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search	Date of mailing of the international search report
30 DECEMBER 1995	29 JAN 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Form PCT/ISA/210 (second sheet)(July 1992)*

Authorized officer

KERRY D. HENDRICKS

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/11855

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Proceedings of the National Academy of Science, USA, Volume 86, issued March 1989, Gould et al., "Use of the DNA polymerase chain reaction for homology probing: Isolation of partial cDNA or genomic clones encoding iron-sulfur protein of succinate dehydrogenase from several species", pages 1934-1938, see entire document.	1-21

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/11855

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☒ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1))(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/11855

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-12, 14-19 and 21 drawn to DNA encoding fatty acid hydroxylase, plants containing the gene, method of using and method of making.

Group II, claims 13 and 20, drawn to oil produced by plants.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The groups are unrelated both physically and chemically, and each possess separate industrial applicability and modes of operation. Further, it is set forth in Annex B of the PCT Administrative Instructions that the method for determining unity of invention under Rule 13 shall be construed as permitting, in particular, the inclusion of the following combination of claims of different categories in the same application:

(i) in addition to an independent claim for a given product, an independent claim for a process specifically adapted for the manufacture of the said product, and an independent claim for a use of the said product.

